

## Identifying slab-derived volatile contributions and mantle source heterogeneity beneath the Washington Cascades

*Kristina Walowski (Western Washington University; walowski@wwu.edu)*

*Paul Wallace (University of Oregon; pwallace@uoregon.edu)*

*Susan DeBari (Western Washington University; susan.debari@wwu.edu)*

*James Rea (University of Oregon; jrea333@gmail.com)*

Fluid-flux melting is commonly attributed as the primary driver of magmatism in arcs globally; however, decompression melting, mantle heterogeneity, and the amount and compositions of slab-derived materials (fluids vs. melts; crust vs. sediments), have all been suggested to play a role in producing the compositional diversity of primary arc basalts. In this contribution, we present the volatile, major, and trace element compositions of melt inclusions hosted from basaltic magmas erupted at three volcanic centers in the Washington Cascades: Mount St. Helens (2 basaltic tephra from the Castle Creek period, 2.0–1.7 ka), Indian Heaven Volcanic Field (two <600ka basaltic hyaloclastitic tuffs), and Glacier Peak (basaltic tephra from Whitechuck and Indian Pass cones). Melt inclusions have H<sub>2</sub>O and Cl contents that range from 0.32–2.15 wt% and 97–1011 ppm, respectively, and are hosted in high Fo olivine (Fo<sub>74</sub>–Fo<sub>85</sub>, Fo<sub>80</sub>–Fo<sub>87</sub>, and Fo<sub>86</sub>–Fo<sub>89</sub>, from Mt St. Helens, Indian Heaven, and Glacier Peak, respectively). We find that trace element ratios, such as Nb/Y, demonstrate that the variability between samples is likely related to differences in mantle source compositions. Impressively, these ratios span nearly the entire range of arc magmas globally, from high Nb/Y compositions at Mount St. Helens that are similar to ocean island basalts, to low-K tholeiites from Indian Heaven and Glacier Peak that have Nb/Y ratios similar to N-MORB. Interestingly, a calc-alkaline basalt from Glacier Peak displays S/Y ratios that overlap with the highest measured in the global arc array from Ruscitto et al. (2012). While all magma types, including calc-alkaline magmas from Indian Heaven and Glacier Peak, have H<sub>2</sub>O and Cl contributions from the down-going plate (inferred from H<sub>2</sub>O/Y and Cl/Y ratios) that overlap with other Cascade Arc segments, the maxima measured in the Washington Cascades are markedly lower than those from other segments, including central Oregon, northern California, and the northern Garibaldi belt, consistent with Venugopal et al. (2020). This dataset adds to the growing inventory of primitive magma volatile concentrations along the Cascade Arc and provides insight into spatial distributions of mantle heterogeneity and the variable role of slab processes in the petrogenesis of arc magmas.

