

A new catalogue of historical Korean auroral records during 1012-1811

Yuqi Wang^{1,2†}, Si Chen^{1,2†}, Kaihua Xu^{1,2†}, Limei Yan^{1,2,3}, Xinan Yue^{1,2,3}, Fei He^{1,2,3*}, Yong
Wei^{1,2,3*}

¹Key Laboratory of Earth and Planetary Physics, Institute of Geology and Geophysics, Chinese
Academy of Sciences, Beijing, China

²College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing,
China

³Beijing National Observatory of Space Environment, Institute of Geology and Geophysics,
Chinese Academy of Sciences, Beijing, China

[†]These authors contribute equally: Yuqi Wang, Si Chen, Kaihua Xu.

*Correspondence to Fei He (hefei@mail.iggcas.ac.cn) and Yong Wei (weiy@mail.iggcas.ac.cn)

Abstract

Aurora displays provides an essential diagnostic to spatial and temporal variations of terrestrial
space environment and is also an important proxy of solar activity. Contemporary auroral
observations have just continued for more than half a century. In the long history prior to modern
era, visual auroral observations can dates back to 1450 AD in mid-latitudes and polar regions in
Europe. In mid- and low-latitude regions in East Asia, official historical books in China, Korea, and
Japan also recorded numerous visual auroral phenomena began from 1000 AD until modern times.
In this study, we compiled a new auroral catalogue from ancient Korean historical books, including

23 2013 auroral records with day-level resolution from 1012 to 1811 AD, especially for the records
24 searched from the *Seungjeongweon Ilgi*. The occurrence of the aurora in the new catalogue is
25 generally consistent with previous datasets. This extended dataset provides valuable support for
26 various studies related to solar-terrestrial space weather and ancient climates.

27 **Key Points:**

- 28 1. Compilation of a new catalogue of historical Korean auroral records during 1012-1811.
- 29 2. The dataset contains 2013 red auroral records with date, local time and direction information.
- 30 3. The dataset provides valuable support for various studies related to solar-terrestrial space weather
31 and ancient climates.

32

33

34 1. Introduction

35 Since Fritz (1873) first identified an ancient aurora list from historical books, many astronomers
36 and space physicists have constructed regional or global lists of ancient auroral records for different
37 purposes with respective emphasizes, such as Link (1964) for years before 1700, Silverman &
38 Blanchard (1983) for England observations, Lee et al. (2004) from Korea histories, and Hayakawa
39 et al. (2017a) for Chinese ancient results. Based on these auroral lists, a variety of studies have been
40 done over decades, especially on estimating the ancient solar and geomagnetic activities given their
41 dominated effect on the aurora occurrence. For example, the auroral records from the Orient played
42 an important role in confirmation of the Maunder Minimum of solar activity (Eddy, 1976), Siscoe
43 (1980) investigated global aurora occurrence based on five available aurora catalogues and
44 identified most known grand solar minimums covering more than 2000 years. He also resolved 80-
45 and 10-year cycle during the Middle Ages, which implies that the solar cycle was then operative.
46 Silverman (1992) analyzed 45000 visual aurora observations during the past 500 years, examined
47 its secular variation through comparison with other solar and magnetic indices, and confirmed the
48 disappearance of 11-year cycle around the Maunder minimum. González-Esparza &
49 Vuelas-Carnoda (2018) used the naked eye aurora observations from Mexico to determine the
50 amplitude of the 1859 great geomagnetic storm and storm evolution, which actually makes
51 significant sense on the extreme storm occurrence estimation for the technique system design given
52 relatively shorter modern observations are available. Furthermore, ancient auroras are also used to
53 investigate the geomagnetic field evolution (Siscoe & Verosub, 2012) and even assist
54 archaeomagnetic dating (Liritzis, 1988)⁰.

55 In these available ancient aurora lists, some of them actually have shown that Korean ancient books

56 recorded a large number of ancient auroras, especially around the Maunder minimum (Dai & Chen,
57 1980). However, due to the relatively low geomagnetic latitude of Korea Peninsula, the current
58 aurora borealis theory cannot explain a large number of auroral records especially in the year of
59 extremely low solar activity. Therefore, in previous studies, most researchers selectively ignored the
60 auroral records of ancient Korea. Here, we systematically collated the auroral records from
61 *Koryo-Sa* (AD 918–1391), *Choson Wangjo Sillok* (AD 1392–1910) and *Seungjeongweon Ilgi* (AD
62 1623–1910) through formatted digital search. Then we manually checked each entry from original
63 books to get detailed information of azimuth, date, local time, lasting time, and etc. Finally, we got
64 2013 auroral records during 1012–1811 and formed this auroral catalogue. In comparison with
65 previous results, this Korean auroral catalogue has several outstanding advantages. Firstly, it was
66 formed based on daily record, which implies that it has time resolution up to one day. Secondly, it
67 was observed from a fixed location with azimuth and local time provided. Thirdly, it has plenty of
68 observations around the Maunder minimum. We expect and believe that the newly created Korean
69 auroral catalogue could be used in a variety of researches in the future.

70 **2. Data and Method**

71 **2.1 Auroral records in Korean chronicles**

72 The Korean chronicles are daily official records of the activities of the kings, the state affairs, and
73 the weather and astronomical phenomena. The existing historical records began in the 1000s and
74 lasted for more than 800 years. Three Korean official historical books: *Koryo-Sa* (AD 918–1391, or
75 *History of Koryo*), *Choson Wangjo Sillok* (AD 1392–1910, or *The Veritable Records of the Choson*
76 *Dynasty*) and *Seungjeongweon Ilgi* (AD 1623–1910, or *The Daily Records of the Royal Secretariat*
77 *of Joseon Dynasty*) were used in this study. These extensive chronicles were written in Chinese

78 characters.

79 The most frequently recorded nocturnal sky glows in the above three books are “vapours like fire
80 light” (pronunciation in Chinese: *qì rú huǒ guāng*) (Stephenson & Willis, 2008). In Chinese, fire
81 light (*huǒ guāng*) refers to red light. This implies that these records represent red glows. In previous
82 works, the records “vapours like fire light” are processed as auroras. Dai and Chen (1980) first
83 systematically sorted out the auroral records in historical books of China, Korea and Japan, but
84 focused on the northern light (auroral borealis), meaning the “vapours like fire light” occurred at
85 north of Korea. Then, Zhang (1985) preliminarily interpreted these records as stable auroral red arcs.
86 Conjugate observations definitely reveal that such “vapours like fire light” in the northern nocturnal
87 sky are auroral borealis (Willis et al., 1996; Hayakawa et al., 2017b). Yau et al. (1995) reorganized
88 the auroral records sorted in previous works (Stephenson & Willis, 2008; Keimatsu, 1970-1976)
89 and published the first comprehensive catalogue of auroral records in East Asian in English.
90 Comparing auroral records in China and Japan with the geomagnetic activity in the 19th century,
91 Willis et al. (2007) propose that the airglow phenomena in the middle and low latitudes is sporadic
92 auroras, as those observed in the United States.

93 Figure 1 shows a drawing of the auroral borealis in historical book of Japan in AD 1770. During
94 this event, although auroral records are not found in *Seungjeongweon Ilgi*, possibly due to the bad
95 weather in Korea⁰, it is believed that the records of “vapours like fire light” in Korean documents
96 refer to the similar auroral structures in Figure 1. Apart from the records in the north, most of the
97 records of “vapours like fire light”, “red vapours” (pronunciation in Chinese: *chì qì*) and “odd red
98 vapours” (pronunciation in Chinese: *chì jìn*) appeared in the southern nocturnal sky. Recently, Wei
99 and Wan (2020) compiled a new chronology of such auroral records in Chinese from the
100 above-mentioned historical books, including 2013 records during AD 1012-1811.





Figure 1. A drawing of auroral borealis in historical book of Japan in AD 1770 (adapted from Hayakawa et al. (2017)).

2.2 Compilation of the dataset

In history, the Korean Peninsula and the Chinese dynasties maintained close astronomical communications, and they both followed a unified standard of astronomical observations and judgement. The red aurorae were depicted by the Chinese character “*qì* (气)” with different adjectives, including “*chì qì* (赤气)” (red vapours), “*qì rú huǒ* (气如火)”/“*qì rú huǒ guāng* (气如火光)” (vapours like fire or vapours like fire light), “*rú huǒ qì* (如火气)” (fire-like vapours), and “*chì jìn* (赤衽)” (odd red vapours). These keywords are used to search the auroral records in the three Korean official historical books.

The National Institute of Korean History has digitalized the three books, and we can quickly search the interested contents using the above keywords from the books’ websites. Each pages of the books are also scanned and labelled with a unique identification number (IDN, e.g., IDN=SJW-A24020230-00200 for the case shown in Figure 2).



116

117 **Figure 2.** An example of Korean red equatorial auroral record. (a) Part of a page from the
 118 Sunjongwon Ilgi containing a red airglow observation. The scanned copy of this record can be
 119 accessed at the National Institute of Korean History
 120 (<http://sjw.history.go.kr/id/SJW-A24020230-00200>). (b) Definition of local directions. (c)
 121 Definition of local times.

122 First, search the three books with the above five keywords to establish a preliminary dataset, in
 123 which there may be false/dummy records or records with incomplete information (date or time).
 124 Therefore, it is necessary to set up a series of criteria to eliminate the bad records. The criteria are as
 125 follows:

- 126 - The record contains one of the keywords and should at least contain date information. Most of
 127 the records contain the full information of date, local time, and observed directions.

128 - The phenomena should be observed at night, i.e. 19:00 LT to 05:00 LT, to eliminate the
129 influence of twilight, solar halo, and other light sources such as clouds, planets, stars, comets,
130 and meteors.

131 - If fire disaster is mentioned in that day, the record is excluded.

132 - Since there is overlap in time for the three books, records from different books on the same day
133 are merged as one and the IDN's for each book are kept in the dataset.

134 After application of these criteria, the final dataset contains 2013 ancient Korean auroral records.

135 Figure 2 shows an example from the *Seungjeongweon Ilgi*. This observation was made in the third
136 year of Emperor Shunzi (the 3rd emperor of the Qing Dynasty), in the 2nd lunar month, and on the
137 23rd day as shown in the 1st and 2nd red rectangles in Figure 2a. the lunar month and dates were
138 converted to the Gregorian calendar with the Buddhist Studies Time Authority Databases
139 (<https://authority.dila.edu.tw/time/index.php>), and the description “*shùn zì sān nián bǐng xū èr yuè*
140 *èr shí sān rì*” is equivalent to April 8, 1646. A translation of the red auroral record is highlighted by
141 the 3rd-5th red rectangles. Their Chinese pronunciation are shown in the square brackets,
142 respectively.

143 For local directions shown in Figure 2b, there are primarily eight directions of *kǎn*, *gèn*, *zhèn*, *xùn*,
144 *lí*, *kūn*, *duì*, and *qián* in ancient China, also called “eight trigrams”, corresponding to north (N),
145 northeast (NE), east (E), southeast (SE), south (S), southwest (SW), west (W), and northwest (NW),
146 respectively. It is noted that the four directions NE, SE, SW and NW are also spelled in Chinese as
147 “*dōng běi*”, “*dōng nán*”, “*xī nán*” and “*xī běi*”, respectively, in ancient documents. Denotations such
148 as “*dōng běi nán*” in the record should refer to three different directions of E, N, and S, meaning
149 that the aurora occurred simultaneously in the three directions.

150 For local time, one day was divided into twelve regular divisions in ancient China, and their
151 correspondences to local times are shown in Figure 2c. Particularly, the nighttime after sunset and
152 before sunrise was divided into five *gēng*'s or five watches, i.e., *yī gēng* (first watch, 19–21 h), *èr*
153 *gēng* (second watch, 21–23h), *sān gēng* (third watch, 23–1 h), *sì gēng* (fourth watch, 1–3 h), and *wǔ*
154 *gēng* (fifth watch, 3–5 h).

155 3. Results and Discussion

156 The ancient auroral dataset presented in this paper was compiled from three Korean official
157 historical books: *Koryo-Sa*, *Choson Wangjo Sillok* and *Seungjeongweon Ilgi*. The dataset is
158 deposited as an Excel table (Ancient Korean Aurora.xlsx) and the corresponding scanned copies of
159 the original books are deposited as PDF files named after the IDN of each auroral record (IDN.pdf).
160 The table contains 8 columns. The 1st column showed the index of the record, the 2nd-4th columns
161 stored the calendar year, month and date, which were converted to the Gregorian date, the 5th and
162 6th columns stored the lunar month and lunar date, the 7th column stored the description of the
163 auroral record translated from the original books, and the 8th column stored the IDN, through which
164 the scanned copy of the original texts can be accessed. One can also find the scanned copy of the
165 auroral records from the three Korean official historical books using the following links.

166 *Koryo-Sa*: <http://db.history.go.kr/id/IDN> (e.g., IDN=kr_053_0010_0030_0100_1000)

167 *Choson Wangjo Sillok*: <http://sillok.history.go.kr/id/IDN> (e.g., IDN=waa_10201025_001)

168 *Seungjeongweon Ilgi*: <http://sjw.history.go.kr/id/IDN> (e.g., SJW-C14020201-02000)

169 Totally, there are 192 records found from *Koryo-Sa*, 536 records from *Choson Wangjo Sillok*, and
170 1359 records from *Seungjeongweon Ilgi*. After merging duplicate records from either book pair, the
171 final number of the auroral records is 2013. All the original Chinese texts are downloaded from the

172 website of the National Institute of Korean History and deposited to the World Data Center for
173 Geophysics, Beijing, together with the Excel table.

174 In 2004, Lee et al. (2004) published 788 auroral records (containing one duplicated record on
175 1537.6.13) from five historical documents of Korea in the 11th–18th century. Most of the records
176 are collected from the three books used in this paper. However, many records are missing in Lee’s
177 list due to manual search of the documents and the original Chinese texts are not appended.
178 Benefitted greatly from the digitalization of the three books by the National Institute of Korean
179 History, systematic and complete search can be done to compile a more comprehensive dataset.

180 In compilation of the dataset, the auroral records are limited to the descriptions of “*chì qì*” (red
181 vapours), “*qì rú huǒ*”/“*qì rú huǒ guāng*” (vapours like fire or vapours like fire light), “*rú huǒ qì*”
182 (fire-like vapours) and “*chì jìn*” (odd red vapours), and four criteria are applied to exclude false/bad
183 records. Figure 3 depicts the distributions of the 2013 ancient Korean auroral records. Figure 3a
184 demonstrates that the majority of the auroral records occur between 1500 and 1800. There are
185 scarce records before 1500 and the distribution is in consistent with Lee’s list (Lee et al., 2004).
186 After 1500, more records are found from *Seungjeongweon Ilgi*. Figure 3b shows the seasonal
187 variation of the records. A peak of the occurrence of the aurora appears in the March, which is
188 consistent with the result of Stephenson’s work (Stephenson & Willis, 2008). Figure 3c presents the
189 variation of the records with lunar date. It is found that there are fewer records around full moon
190 and more records around new moon.




Figure 3. Statistics on the distributions of the ancient Korean auroral records. (a) Histogram of auroral records versus year binned in 10-year intervals. (b) Histogram of auroral records versus month. (c) Histogram of auroral records versus lunar date binned in 3-day intervals.

4. Summary

In this work, a new catalogue of historical Korean auroral records during 1012-1811. The dataset contains 2013 red auroral records with date, local time and direction information. The compiled

dataset is provided as XLSX file which can be opened in Excel or other text editors. The original Chinese texts are provided as PDF files which can be opened in PDF Reader. All these files have been deposited to the World Data Centre for Geophysics, Beijing (<http://www.geophys.ac.cn/ArticleDataInfo.asp?MetaId=207>, doi: 10.12197/2020GA008) and can be permanently accessed. The dataset can be freely used for research and education purposes only. The dataset provides valuable support for various studies related to solar-terrestrial space weather and ancient climates.

Acknowledgements

We greatly appreciate the National Institute of Korean History for providing the public access of the historical books available at <http://www.history.go.kr>. We thank National Diet Library of Japan for providing the auroral drawing. We also thank the Dharma Drum Institute of Liberal Arts to operate the Buddhist Studies Time Authority Databases (<https://authority.dila.edu.tw/time/index.php>). This work is supported by the Strategic Priority Research Program of Chinese Academy of Sciences (Grant XDA17010201), the National Natural Science Foundation of China (41525016), the Key Research Program of the Institute of Geology & Geophysics, CAS (grant IGGCAS-201904), and the Youth Innovation Promotion Association of the Chinese Academy of Sciences (2017258).

References

- Dai, N. Z. & Chen, M. D. (1980). Chronology of northern light in history of China, Korea and Japan: From the legendary period to 1747 AD. *Kejishiwenji* (in Chinese), **6**, 87–146
- Eddy, J. A. (1976). The Maunder Minimum. *Science*, **192**, 1189-1202.
- Fritz, H. (1873), *Verzeichniss beobachteter Polarlichter*, Wien, Austria.

220 González-Esparza, J. A. & Cuevas-Cardona, M. C. (2018). Observations of low-latitude red aurora
 221 in Mexico during the 1859 Carrington geomagnetic storm. *Space Weather*, **16**, 593-600.

222 Hayakawa, H. et al. Records of sunspots and aurora candidates in the Chinese official histories of
 223 the Yuán and Míng dynasties during 1261–1644. *Publications of the Astronomical Society of*
 224 *Japan*, **69**, 65, doi:10.1093/pasj/psx045 (2017a).

225 Hayakawa, H. et al. (2017). Long-lasting extreme magnetic storm activities in 1770 found in
 226 historical documents. *The Astrophysical Journal Letters*, **850**, L31, doi:
 227 10.3847/2041-8213/aa9661

228 Keimatsu M., 1970. "A chronology of aurorae and sunspots observed in China, Korea and Japan".
 229 *Annals of Science, Kanazawa Univ. Part I*, **7**, 1-10

230 Link, F. (1964). Observations et catalogue des aurores boréales apparues en occident de 1601 a
 231 1700. *Geofysica Sbornik*, **212**, 501–550

232 Lee, E. H. et al. (2004). The Sunspot and Auroral Activity Cycle Derived from Korean Historical
 233 Records of the 11th–18th Century. *Solar Physics*, **224**, 373–386

234 Liritzis, Y. (1988). Aurorae boreales and geomagnetic inclinations as aids to archaeomagnetic dating.
 235 *Earth, Moon, and Planets*, **42**, 151-162

236 Silverman, S. M. & Blanchard, D. (1983). Wilson Bentley's auroral observations. *Planetary and*
 237 *Space Science*, **31**, 1131–1135

238 Silverman, S. M. (1992). Secular variation of the aurora for the past 500 years. *Review of*
 239 *Geophysics*, **30**, 333– 351

240 Siscoe, G. L. (1980). Evidence in the auroral record for secular solar variability. *Review of*
 241 *Geophysics*, **18**, 647– 658

242 Siscoe, G. L. & Verosub, K. L. (2012). High medieval auroral incidence over China and Japan:

243 Implications for the medieval site of the geomagnetic pole. *Geophysical Research Letters*, **10**,
 244 345-348

245 Stephenson, F. R. & Willis, D. M. (2008). ‘Vapours like fire light’ are Korean aurorae. *Astronomy &*
 246 *Geophysics*, **49**(3), 3.34–3.38, doi:10.1111/j.1468-4004.2008.49334.x

247 Wei, Y. & Wan, W. *Chronology of Ancient Korean Aurora* (in Chinese) (Science Press, Beijing,
 248 2020).

249 Willis, D. M., Stephenson, F. R. and Singh, J. R. (1996). Auroral observations on AD 1770
 250 September 16: the earliest known conjugate sightings. *Quarterly Journal of the Royal*
 251 *Astronomical Society*, **37**, 733–742

252 Willis, D. M., Stephenson, F. R. & Fang H. (2007). Sporadic aurorae observed in East Asian.
 253 *Annales Geophysicae*, **25**, 417–436

254 Yau, K. K. C., Stephenson, F. R. & Willis, D. M. (1995), A catalogue of auroral observations from
 255 China, Korea and Japan (193 B.C. – A.D. 1770). Report No. RAL-TR-95-073, Rutherford
 256 Appleton Laboratory, Harwell Oxford.

257 Zhang, Z. W. (1985). Korean auroral records of the period AD 1507-1747 and the SAR arcs.
 258 *Journal of the British Astronomical Association*, **95**(5), 205–210

259