

Fig. 1 Temporal response of net CO₂ assimilation (A) and stomatal conductance (g_s) to a step change in PPFD from 200 (shaded area) to 1000 (unshaded area) $\mu\text{mol m}^{-2} \text{s}^{-1}$ in *G. max*. (a, f) 0 day after treatments (DAT), (b, g) 2 DAT, (c, h) 4 DAT, (d, i) 5 DAT, (e, j) 7–8 DAT. Orange, grey, green, and blue lines indicate control (Ct), cold-girdling (CG), sucrose feeding (Suc), and low nitrogen (LN) treatments, respectively. Values are mean \pm SD with a measurement interval of 10 seconds obtained from Exp. 1–3 ($n=4\text{--}12$).




Fig. 2 Temporal response of net CO₂ assimilation (A) and stomatal conductance (g_s) to a step change in PPFD from 200 (shaded area) to 1000 (unshaded area) $\mu\text{mol m}^{-2} \text{s}^{-1}$ in *P. vulgaris*. (a, e) 0 day after treatments (DAT), (b, f) 2 DAT, (c, g) 5~6 DAT, (d, h) 7 DAT. Orange, grey, green, and blue lines indicate control (Ct), cold-girdling (CG), sucrose feeding (Suc), and low nitrogen (LN) treatments, respectively. Values are mean \pm SD with a measurement interval of 10 seconds obtained from Exp. 1–2 ($n=4-8$).




Fig. 3 Nonstructural carbohydrate (NSC_{area}) and leaf nitrogen content per area (N_{area}) in the primary leaves of *G. max* and *P. vulgaris* on 7–8 days after treatments (DAT). (a, c) *G. max* and (b, d) *P. vulgaris*. Orange, grey, green, and blue bar indicate control (Ct), cold-girdling (CG), sucrose feeding (Suc), and low nitrogen (LN) treatments, respectively. NSC_{area} is expressed as sum of starch, sucrose, and glucose. Triangle represents data of *G. max* in Exp. 3. Bars are mean + SD obtained from Exp. 1–3 (n=4–12). Different lower-case letters indicate significant differences among treatments (Tukey's test, $P < 0.05$).




Fig. 4 Relationships between photosynthetic traits (A_{\max} and g_{smax}), nonstructural carbohydrate (NSC_{area}), and leaf nitrogen content per area (N_{area}) in the primary leaves of *G. max* on 7–8 days after treatments (DAT). Orange, grey, green, and blue symbols indicate control (Ct), cold-girdling (CG), sucrose feeding (Suc), and low nitrogen (LN) treatments, respectively. Triangle represents data of *G. max* in Exp. 3. Values of R^2 are (a) 0.77, (b) 0.71, (c) 0.86, (d) 0.61 ($P < 0.01$).




Fig. 5 K-means clustering and gene ontology (GO) enrichment analysis from transcriptome data of *G. max* obtained in Exp. 3. Heat map shows up- (red) and down- (green) regulated genes in control (Ct), cold-girdling (CG), and low nitrogen (LN) (n = 4). 2000 genes were grouped into 4 clusters, each of which include 442 (Cluster A), 557 (Cluster B), 473 (Cluster C), and 508 genes (Cluster D) (Supplementary Table X). Top 10 GO biological processes were shown for each cluster.




Fig. 6 Expression of key genes in the Calvin cycle in response to cold-girdling (CG) and low nitrogen (LN). Data are obtained from transcriptome data of *G. max* in Exp. 3. Transcript levels of genes annotated with Arabidopsis gene ID are calculated as the total read counts of duplicated genes of *G. max* (Supplementary Table 3). RCA is Rubisco activase that facilitates Rubisco activation, and RAF is Rubisco accumulation factor required for assembly and stability of Rubisco. PRK is responsible for the regeneration of RuBP. Bars are mean \pm SD ($n=4$). Different lower-case letters indicate significant differences among treatments (Tukey's test, $P < 0.05$).




Fig. 8 Changes in expression of gene involved in nitrogen assimilation in response to cold-girdling (CG) and low nitrogen (LN). Data are obtained from transcriptome data of *G. max* in Exp. 3. Transcript levels of genes annotated with Arabidopsis gene ID are calculated as the total read counts of duplicated genes of *G. max* (Supplementary Table 8). NRT and AMT are NO_3^- and NH_4^+ transporter, respectively, where NRT1.7 is responsible for phloem loading of NO_3^- . NR and NiR are nitrate and nitrite reductase, respectively. GLN, FD-GOGAT, and ASN are glutamine (Gln), glutamate (Glu), and asparagine (Asn) synthase, respectively. CAT and AAP are cationic and neutral amino acid transporters, respectively. Bars are mean \pm SD (n=4). Different lower-case letters indicate significant differences among treatments (Tukey's test, $P < 0.05$).




Fig. 9 Changes in expression of gene involved in starch synthesis and sugar transport/metabolism in response to cold-girdling (CG) and low nitrogen (LN). Data are obtained from transcriptome data of *G. max* in Exp. 3. Transcript levels of genes annotated with Arabidopsis gene ID are calculated as the total read counts of duplicated genes of *G. max* (Supplementary Tables 9-11). APL and APS are large and small subunits of ADP glucose pyrophosphorylase (AGPase), respectively. SS and GBSS are soluble and granule bound starch synthases, respectively. TPT and GLT export triose phosphate (TP) and glucose from the chloroplast to the cytosol, respectively. GPT transport glucose-6-phosphate (G6P) from the cytosol to the chloroplast. SPS and SPP play a major role in sucrose synthesis, and SUT is sucrose transporter. Bars are mean \pm SD (n=4). Different lower-case letters indicate significant differences among treatments (Tukey's test, $P < 0.05$).