

Assessment of diurnal urban heat island (UHI) intensity in microclimatic urban environment using Local climate zone classification approach


Session Number : GC33G-1225 Abstract ID: 1375839

Ashish Mishra*, Dhyans Arya, Department of Hydrology, Indian Institute of Technology Roorkee, India

ABSTRACT


- Adoption of Local Climate Zone (LCZ) classification to study UHI in a microclimatic urban setting, focusing on Dehradun city's diverse urban landscape.
- Combined use of remote sensing data and ground-based measurements to evaluate the temporal and spatial variations of UHI intensity, both during day and night.
- Discovery of distinct diurnal patterns in UHI intensity, with peak occurrences in the late afternoon and early evening across different LCZs.

MOTIVATION



Source: Kang S et.al. (2022)


STUDY AREA AND METHODOLOGY



- ❑ The municipal region of the city, covering an area of approximately 74 sq. km.
- ❑ Average annual temperature of 20.5 °C
- ❑ Warm and temperate climate
- ❑ Period of Analysis March 2020 to November 2022.



Weather Station information in Dehradun city




Height of
Instrument: 2 m
Location : Flat
ground (A, D) and
Building rooftop
(B,C)

Data: Sub-hourly interval (15 min)- **Temperature, Relative Humidity, Wind Speed, Wind Direction, Rainfall, Dew Point, Solar Radiation**

$$UHI \text{ Intensity } (^{\circ}C) = T_{urban} - T_{rural}$$

KEY FINDINGS

- Identification of diverse LCZs in Dehradun, including compact low-rise, mid-rise areas, dense trees, and open spaces.
- Discovery of distinct diurnal UHI patterns with peak intensities during late afternoon and early evening.
- Notable impact of vegetation and built-up areas on UHI variation – cooling effect of green spaces and amplifying effect of impervious surfaces.
- The rural station AWS_78 was used to compute the UHI intensity for all urban stations. The mean UHI intensity by hour of the day was plotted for all station.
- AWS 75, AWS 77, and AWS 80 exhibit a pronounced diurnal pattern with higher UHI intensity during the late afternoon and early evening hours, typically peaking between 1500 to 2000 hours.



LCZ Classification map for Dehradun city

- AWS 76 and AWS 79 show a more uniform distribution of UHI intensity throughout the day, with no significant peaks indicating a less pronounced diurnal UHI effect.

CONCLUSIONS

- ❑ Compact urban areas exhibit more stable UHI intensities, while open low-rise zones show greater diurnal variation.
- ❑ The intensity of UHI peaks in areas with lower-rise buildings, suggesting a significant impact of urban morphology on UHI.
- ❑ The variation of UHI across LCZs highlights the critical role of built form in urban thermal dynamics.

ACKNOWLEDGEMENT

The authors gratefully acknowledge the Indian Institute of Technology Roorkee, India for funding and resources, the AGU for the Student Travel Grant, and the Department of Science and Technology (DST-SERB), Government of India for the International Travel Grant (ITS/2023/005179), enabling attendance at the AGU Fall Meeting 2023.

CONTACT INFORMATION



Ashish Mishra



Dhyans Arya

