

# Biomechanical modelling of cancer: Agent-based force-based models of solid tumours within the context of the tumour microenvironment

Cicely K Macnamara<sup>a,\*</sup>

<sup>a</sup>*School of Mathematics and Statistics, Mathematical Institute, University of St Andrews, United Kingdom, KY16 9SS*

---

## Abstract

Once cancer is initiated, with normal cells mutated into malignant ones, a solid tumour grows, develops and spreads within its microenvironment invading the local tissue; the disease progresses and the cancer cells migrate around the body leading to metastasis, the formation of distant secondary tumours. Interactions between the tumour and its microenvironment drive this cascade of events which have devastating, if not fatal, consequences for the human host/patient. Among these interactions, biomechanical interactions are a vital component. In this paper, key biomechanical relationships are discussed through a review of modelling efforts by the mathematical and computational oncology community. The main focus is directed, naturally, towards lattice-free agent-based, force-based models of solid tumour growth and development. In such models interactions between pairs of cancer cells (as well as between cells and other structures of the tumour microenvironment) are governed by forces. These forces are ones of repulsion and adhesion, and are typically modelled via either an extended Hertz model of contact mechanics or using Johnson-Kendal-Roberts theory, both of which are discussed here. The role of the extracellular matrix in determining disease progression is outlined along with important cell-vessel interactions which combined together account for a great proportion of Hanahan and Weinberg's *Hallmarks of Cancer* [1, 2].

*Keywords:* agent-based; force-based; in silico tumours; cancer growth and development; tumour microenvironment

---

## 1. Introduction

The term cancer covers a spectrum of diseases – cancer cells can arise from any type of cell in the body and can grow in or around any tissue or organ making it highly complex. Tumour cells proliferate, occupying whole areas of

---

\*Corresponding author

Email address: [ckm@st-andrews.ac.uk](mailto:ckm@st-andrews.ac.uk) (Cicely K Macnamara)

5 tissue; they interact with surrounding cells, tissue structures, vasculature and  
 6 the extracellular matrix (ECM) in a variety of ways. In recent years, mathem-  
 7 atical and computational biologists have endeavoured to accurately capture the  
 8 growth and development of tumours within their local environment through *in*  
 9 *silico* models. By simulating virtual tumours, insight is gleaned which comple-  
 10 ments traditional biological and experimental approaches to cancer research at  
 11 limited financial and ethical cost. This paper will focus on highlighting selec-  
 12 ted lattice-free agent-based (specifically force-based models) of tumour growth  
 13 and development. By way of introduction it will be worthwhile to discuss the  
 14 importance of approaching the problem from a mechanical standpoint, as such,  
 15 in Section 1.1, the tumour microenvironment (TM) is presented followed by, in  
 16 Section 1.2, a discussion of the inherent biomechanics of the TM. In Section 1.3  
 17 certain other modelling techniques which have been used to study the dynamics  
 18 of tumour growth and development will be highlighted paying specific attention  
 19 to where biomechanics have been successfully implemented.

### 20 1.1. The tumour microenvironment (TM)

21 The term *tumour microenvironment* is given to all aspects of the local en-  
 22 vironment of a tumour, consisting of, but not limited to, the surrounding blood  
 23 vessels/vasculature, ECM, tumour-associated immune cells and signalling mo-  
 24 lecules/proteins released by the cancer cells (see schematic in Figure 1). The  
 25 tumour and the TM are intrinsically linked and there is constant interplay and  
 26 interactions between them starting from the point of tumour initiation [3]. In-  
 27 deed, non-cancerous cells within tissue respond continuously to the external  
 28 signals of their environment, changing their metabolic state, growth, mitosis,  
 29 gene expression, differentiation, movement, or even undergoing programmed  
 30 cell death (apoptosis), accordingly. Should the cell fail to correctly transduce or  
 31 respond to a specific (external) signal it effectively becomes cancerous. A cell  
 32 with a cancerous phenotype has several distinct *Hallmarks* [1, 2]. For example,  
 33 cancer cells resist apoptosis and enable replicative immortality; this unchecked  
 34 proliferation creates a tumour (or neoplasm) within the tissue.

35 Tumours influence the TM in a variety of different ways. Hypoxic tumour  
 36 cells, starved of oxygen, are known to release vascular-endothelial growth factor  
 37 (VEGF) which promotes tumour angiogenesis, supplying the tumour with con-  
 38 stant access to vital nutrient [4, 5]. Equally, as the growing tumour vies for  
 39 space within the tissue, cells release matrix metalloproteinases (MMPs) which  
 40 degrade the ECM making room for tumour growth and local invasion [6, 7].  
 41 Conversely, the TM affects tumour growth and development; the shape and size  
 42 of a tumour; but also its genetic evolution being determined by properties of  
 43 the local environment. For example, cells migrate preferentially up gradients of  
 44 ECM stiffness in a specific type of mechanotaxis called *durotaxis* [8]. Stiff ECMs  
 45 can promote tumourigenesis through integrin-dependent mechanotransduction  
 46 at focal adhesions [9] while soft ECMs contribute to phenotypic selection of  
 47 tumour-repopulating cells (TRCs) [10]. Indeed the TM has been found to play  
 48 an active role in the progression of malignancies [11, 12].




Figure 1: Schematic diagram showing several key aspects of the TM: the cancer cells (blue), the ECM fibres (black), the vasculature (red), vascular-endothelial growth factor (VEGF) signalling protein (magenta) and immune cells (orange).

One of the *Hallmarks of Cancer* is tissue invasion and metastasis in which tumours spread both locally and non-locally [1, 2]. Malignant tumours aggressively take over large areas of tissue, and, of greater concern, are able to move from primary locations to secondary locations using the body’s circulatory system. This is a major issue since it is commonly purported that as many as 90% of all cancer deaths are due to metastatic spread; note that this figure while widely reported and hypothesised is not yet scientifically proven although it is true that the majority of cancer fatalities are due to metastases [13]. Nevertheless, agent-based models of tumours typically and vitally should also include aspects of the TM in order to model how cancers invade and metastasise.

### 1.2. Biomechanics in the TM

The focus of this paper is towards force-based models, and as such it is important to understand why mechanical interactions are so important. As discussed above there is constant interplay between a tumour and the TM. Indeed, the TM governs how a tumour establishes and develops; the tumour cells respond to mechanical cues actively by changing shape, state or migrating. For example, Friedl and co-workers have shown how the specific nature of the ECM (it’s density, stiffness and geometry) along with aspects of the cancer cell (it’s adhesive properties and polarity) determine how a cell (or a collection of cells) migrates through tissue [14–18]. *Durotaxis* was mentioned above but another type of “taxis” experienced by cells is *haptotaxis* [19] which is motility of cells preferentially up gradients of adhesion within the ECM. More generally cells are affected by “mechanotransduction”, in which cell-external mechanical stresses provoke cell-internal chemical signals leading to some type of adaptive response [20]. For further discussion of mechanotransduction in cancer see the

review of the same name [21]. Equally, within the tumour itself stresses affect development. Homeostatic pressure in which a balance of proliferation and apoptosis results in zero net growth has been found to limit the growth of some solid tumours [22, 23]. Conversely, such mechanical compression (solid stress) may actually drive cancer cells to invade and metastasise [24–27]. Given the intrinsic links between cancer cell behaviour and biomechanics, in order to fully understand how tumours, initiate, grow, invade and metastasise it is vital to include such processes in mathematical and computational models.

### 1.3. Other *In Silico* models

Early mathematical modelling of cancer (avascular solid tumours) focused on deterministic or continuum models of solid tumour spheroids developed from the classical Greenspan model [28]. Such models continue to provide insight through the ability to efficiently model large scale dynamics (typical palpable tumours will contain at least  $10^8$  cells [29]) and equally since they lend themselves to mathematical analysis. For reviews of deterministic and continuum models see, for example, [30, 31]. Selected articles in which mechanical stress is modelled using a continuum approach include [3, 32–36] while cell-cell interactions are considered in [37–43], and cell-matrix interactions in [44].

More recently efforts have been focused on using individual-based models or agent-based models which allow a more direct comparison to the biology through the ability to model at the cell scale and within. In fact, modelling cell behaviour on the individual level is naturally scale bridging allowing at once intracellular (microscopic) and intercellular (mesoscopic) mechanisms to be included even when modelling a large number of cells (macroscopic). Equally, taking an individual approach easily allows the modelling of heterogeneous cell populations or, at the very least, variability between cells.

#### 1.3.1. On lattice models

The most simplistic agent-based models are cellular automata models; in general, on-lattice agent based models have dominated the literature, these can be broadly categorised into four distinct types (see Table 1). Note, in the schematics in Table 1 each type is shown on a structured square lattice, however, on-lattice models often now use unstructured lattices such as the Voronoi-Delaunay lattice, for example, which typically results in more biologically realistic shapes, both of cells and cell-masses [46, 69]. On-lattice models may be 3D as in the case of the classic multicellular tumour spheroid (MCTS) models or 2D as in the case of monolayers. On-lattice models lend themselves to efficient large scale simulations of a great number of cells at little computational cost. Table 1 provides details of some selected references for state-of-the-art on-lattice models of tumour growth, specifying the tumour-TM interactions considered where appropriate. For further discussion of on-lattice models see, for example, the reviews in [69–75]. On-lattice models typically do not include mechanics which may be necessary to accurately depict the biology (see discussion above). Types I, II and IV rely solely on stochastic processes governing changes of state or position







Schematic	Model Description	Selected References
	Type I - Single cell per lattice site	MCTS [45–47]; cell-vessel interactions [48]; cell adhesion [49–52]; monolayers [53]; phenotypic heterogeneity [54–57]
	Type II - Compartment model. Multiple cells per lattice site	coarse-grained proliferative rim [58, 59]
	Type III - Single cell covers multiple lattice sites (Cellular-Potts)	MCTS [60]; cell adhesion [61, 62]; angiogenesis [63]; cell-fibre interaction [64]; monolayers [65]
	Type IV - Multiple (or single) cell(s) per lattice site, movement through velocity channels (lattice gas cellular automata)	MCTS [66]; cell-fibre interaction [67]; cell-ECM interaction [68]

Table 1: Summary of on-lattice models with some selected references.

of a cell as well as mitosis. Cellular-Potts (Type III) is the only type to permit the modelling of physical mechanisms by solving an effective energy equation which goes some way to modelling the forces between cells (see, for example, [76–78]). There are several open-source on-lattice computational frameworks which include, notably for cancer, the *CompuCell3D* Cellular-Potts framework [79].

The remainder of this paper considers lattice-free (or off-lattice) agent-based, specifically, centre-based, force-based models of tumour growth and development. It is structured as follows: in Section 2 the modelling approach is introduced, in Section 3 the specifics of the forces acting between cells are outlined and in Section 4 there is a discussion of selected modelling efforts of other aspects of the TM. Throughout, a sample of results from the literature will be given.

## 2. Centre-based force-based modelling

Within a lattice-free agent-based model each component (e.g. cell, tissue fibre or vessel segment) is considered explicitly. Let us start by considering the most important aspect of the TM, the tumour cells themselves. Each cancer cell,  $i$ , is an individual agent; this paper focuses on centre-based models (CBM) in which the cell geometry is simplified with each cell considered to be a viscoelastic sphere subject to small deformations, described by the position of its centre,  $\underline{x}_i$ , in the domain (hence centre-based) and its radius,  $R_i$ , see leftmost image of Figure 2. When growing tumours of significant size it is a reasonable assumption/simplification to make that cells may be represented by spheres. Other tumour models exist in which cells have non-spherical shape or are fully deformable, notably the work of Rejniak and coworkers [80–83]). However, these are not the subject of the review given here.




Figure 2: Schematic diagram indicating the basic physical properties of cells in centre-based models, showing on the left a single cell in isolation primed for mitosis, in the middle that seed cell having undergone mitosis creating two daughter cells and on the right two mature cells in contact under a balance of forces.

The behaviour of tumour cells can be broken down into three distinct but linked aspects. Firstly, there are biological factors such as the cell cycle; each cell has the ability to grow in size and divide, undergoing mitosis. Once a cell has reached maturity (proliferative size) it may split into two daughter cells; mitosis is considered a stochastic event (taking place randomly, indicated by the DNA segments on the growth timescale in Figure 2) with probability inverse to the cell-cycle time. When the mother cell divides the simplest implementation is to have two smaller (volume preserving) daughter cells replace the mother cell (see middle image of Figure 2) [84, 85], more sophisticated models depict the splitting more accurately by deforming the spherical mother cell into a dumbbell shape the ends of which eventually separate into the daughter cells [86, 87]. The daughter cells then grow according to a growth rate until they too reach

proliferative size and experience forces imposed by each other (see below and Section 3). Mitosis may be inhibited by external factors such as an excessive compression force due to a high number of neighbouring cells, this is known as contact inhibition [86].

Secondly, there are genetic factors; cells may have given phenotypes or genotypes which prescribe their behaviour in some way. For example, cell phenotypic evolution might depend on biophysical processes, or biochemical interactions such as the availability of nutrients. This will be discussed further in Section 4.3.2, in which the traits of cells with a hypoxic phenotype are compared to the *Hallmarks of Cancer*.

Lastly, and particularly key, for force-based models interactions between cells (and indeed other agents in the model) are described by forces or potentials. Typically, each cell is governed by an equation of motion, an ordinary differential equation of the form:

$$\underbrace{\mathbf{\Gamma}\dot{\mathbf{x}}_i(t)}_{\text{friction}} + \underbrace{f_i(t)}_{\text{migration}} = \underbrace{\sum \mathbf{F}_i(t)}_{\text{mechanical forces}} . \quad (1)$$

The equation of motion takes into account three main aspects. Firstly, it accounts for friction experienced by the cell (first term in Equation (1), in which  $\mathbf{\Gamma}$  is a 3-dimensional tensor that models the physical structure of the environment) - this may be simply background friction imposed by the tissue but may account for friction imposed on cells by other structures. Secondly, the cell will have some pre-described active migration properties (second term in Equation (1)), these may be as simple as random fluctuations/motion as in [84, 85] or may take into account a cells preferred direction (polarity) as in [88] and even effects of the external environment (e.g. chemotaxis where cells are naturally driven up gradients of nutrient, as in [89]). Thirdly, it incorporates mechanical interactions via forces (third term in Equation (1)) between a cell and other agents within the model.

For two cells in contact (determined when the distance between their centres is less than the sum of their radii) a force directed along the vector between their centres,  $\mathbf{d}_{ij}$ , is calculated taking into account repulsion and adhesion. Resolving the resulting potential between the two cells in the absence of any migration terms leads to two cells which remain stationary under a balance of forces (see rightmost image of Figure 2). In the following Section we discuss in more detail the repulsion and adhesion forces between cells. Later we will outline interactions of cells with other aspects of the TM (Section 4).

### 3. Repulsion and Adhesion Forces

Force-based models are naturally governed by forces, specifically, repulsion and adhesion forces. In this Section the repulsion and adhesion forces acting between cancer cells are elucidated. The types of model discussed assume that a cell is spherical in isolation. Thus, any large contact area between a pair of cells (and indeed multiple contact areas between a cell and multiple others)

creates a significant stress on the cytoskeleton of the cell(s). The limited ability to deform or indeed compress (with Poisson numbers found by experiments to be between approximately 0.4 – 0.5 [90]) leads to repulsion between cells. Conversely, cells are naturally adhesive. For cells in contact, binding due to adhesive molecules occurs; as the contact area increases so too do the adhesive bonds. The adhesive molecules at play are Cadherins (calcium-dependent adhesion) and Catenins, together these proteins form complexes called adherens junctions which facilitate cell-cell adhesion. Ramis-Conde and coauthors incorporated the E-Cadherin- $\beta$ -Catenin pathway explicitly into their individual based model of tumour development in order to discuss the implications of this pathway on cell migration and cancer invasion [91–94].

The total cell-cell interaction force between two cells,  $i$  and  $j$ , directed along the vector,  $\mathbf{d}_{ij}$ , joining their centres (see righthmost image of Figure 2), is given by

$$\mathbf{F}_{i,j} = (\mathbf{F}_{i,j}^{\text{rep}} - \mathbf{F}_{i,j}^{\text{adh}}) \frac{\mathbf{d}_{ij}}{\|\mathbf{d}_{ij}\|}, \quad (2)$$

where  $\mathbf{F}_{i,j}^{\text{rep}}$  is the repulsion force discussed in Section 3.1 and  $\mathbf{F}_{i,j}^{\text{adh}}$  is the adhesion force discussed in Section 3.2. In order to calculate the change in position of cell  $i$  at each timestep, the sum of all resulting forces between cell  $i$  and any cell  $j$  with which it is in contact is included in the equation of motion (Equation 1).

### 3.1. Hertzian Repulsion

For two spherical cells,  $i$  and  $j$ , in contact and subject to small (elastic) deformations, the repulsive force experienced is typically described in the literature by the classical Hertzian contact mechanics repulsion [95]. The form of the repulsion force for two such cells of radii  $R_i$  and  $R_j$ , is, therefore

$$|\mathbf{F}_{i,j}^{\text{rep}}| = \frac{4}{3} E^* R^{*1/2} h_{ij}^{3/2}, \quad (3)$$

where  $h_{ij} = R_i + R_j - \|\mathbf{d}_{ij}\|$  describes the length of “overlap” (or contact area) between the two cells. This repulsion force term includes both an effective radius,  $R^* = R_i R_j / (R_i + R_j)$  and an effective Young’s Modulus,  $E^*$ , which is calculated from

$$\frac{1}{E^*} = \frac{1 - \nu_i^2}{E_i} + \frac{1 - \nu_j^2}{E_j}, \quad (4)$$

where  $E_i$  and  $E_j$  are the cells’ respective Young’s moduli and  $\nu_i$  and  $\nu_j$  their Poisson ratios.

Under Hertzian elastic contact alone the following assumptions must be made: (a) strains on the cells are small and within the elastic limit, (b) the area of contact between the spherical cells is much smaller than their radii, (c) the cell surfaces are continuous and non-conforming and (d) there is no friction between the cells. Moreover, this classical model is strictly non-adhesive. Cells, however, are naturally adhesive, governed by adhesion molecules that travel

to the cellular membrane, stimulated by the proximity of a neighbouring cell, forming adhesive bonds. Thus, for those modelling mechanical cell-cell interactions using contact mechanics it is necessary to also include an adhesion force between cells, thus extending or modifying the classical Hertzian model.

### 3.2. Adhesion

There are several examples in the literature of cell-cell interaction forces, with differing expressions for the adhesive force,  $\mathbf{F}_{i,j}^{\text{adh}}$ . Here we discuss two key variants. These are outlined in Table 2 for quick reference and comparison. In each case the force takes into account the strength of adhesion,  $\alpha$ , which is assumed to be constant among the cell population and considers the contact surface area between cells since as contact surface area increases so too does the number of adhesive bonds.

Adhesive Force	Description	References
$ \mathbf{F}_{i,j}^{\text{adh}}  = \alpha S_{ij}$ , e.g. $= 2\pi\alpha \left(R_i - \frac{h_{ij}}{4}\right) h_{ij}$	Adhesion directly proportional to contact surface area, $S_{ij}$ . The resulting force can be determined explicitly.	[84, 85, 91, 92, 96]
$ \mathbf{F}_{i,j}^{\text{adh}}  = \frac{4E^*}{3R^*} a^3 - [8\pi\alpha E^* a^3]^{1/2}$	Johnson-Kendal-Roberts (JKR) theory. Contact surface area (with contact radius parameter $a$ ) is modified by adhesion. The resulting force must be determined implicitly.	[87, 97, 98]

Table 2: Selected forms of CBM adhesion force with selected references.

241

#### 3.2.1. Explicit adhesion force

In this variant, the adhesion force,  $\mathbf{F}_{i,j}^{\text{adh}}$ , between two overlapping cells, is assumed be directly proportional to the contact surface between them,  $S_{ij}$ . The contact surface area is first calculated which then feeds into the adhesion force. Within the literature there are different approximations for the contact surface area. In [96], for example, they model the contact surface area of cells in contact as the area of the circle equidistant between the two cells, underlying the spherical cap of height  $h_{ij}/2$  (i.e. half the overlap between cells). While in [84] they calculate the area to be the average value between the area of the spherical cap of height the overlap between the cells,  $h_{ij}$ , and area of the circle underlying the cap (see Figure 3). In this case, the contact surface is approximated as

$$S_{ij} = \frac{1}{2} [2\pi R_i h_{ij} + \pi (2R_i h_{ij} - h_{ij}^2)] = 2\pi R_i h_{ij} + \frac{\pi h_{ij}^2}{2},$$

253 with the resulting adhesion force given by

$$|\mathbf{F}_{i,j}^{\text{adh}}| = 2\pi\alpha \left( R_i - \frac{h_{ij}}{4} \right) h_{ij}. \quad (5)$$




Figure 3: Figure showing how the contact area is estimated in [84].

254

255

256

257

258

259

260

261

262

263

264

265

266

267

This approach to modelling adhesion considers a “suction” effect as a consequence of the increasing density of effective bonds between the cells. In such an approach certain assumptions have been made [96]. Firstly, it is assumed that the adhesion molecules (receptors and ligands) which bind the cells together are distributed homogeneously over the whole cell surface and thus the whole contact surface area. Secondly, that binding takes place instantaneously and furthermore that since adhesion which causes deformations to the cell naturally change the cell surface area it is assumed that this process happens rapidly so that it is not necessary to explicitly consider the cell surface area.

Figure 4 shows the growth of a MCTS over 3 000 time steps (approximately 2 days) in which adhesion is modelled by the explicit adhesion force given by Equation 5. The simulation results shown are derived from the model (along with parameters) given in [85].

268

### 3.2.2. Implicit JKR adhesion force

269

270

271

272

273

274

The explicit model(s) of adhesion discussed in the previous Section, do not take into account the fact that the adhesion (derived from the surface contact area) then affects and modifies the surface contact area. The Johnson-Kendall-Roberts (JKR) theory of adhesive contact derives a model for the adhesive force which includes this hysteresis phenomena [99]. In this case the force is given by

$$|\mathbf{F}_{i,j}^{\text{adh}}| = \frac{4E^*}{3R^*} a^3 - [8\pi\alpha E^* a^3]^{1/2}, \quad (6)$$

275

276

in which  $E^*$  and  $R^*$  are, once again, the effective Young’s modulus and radius, respectively and  $a$  is the contact surface radius (see Figure 3). However, in this




Figure 4: Figures showing the results of a computational simulation of the growth over time of a MCTS from the CBM of [85] (unpublished result) in which adhesion is incorporated via Equation 5.

case  $a$  is not fixed but rather changes and may be calculated from

$$h_{ij} = \frac{a^2}{R^*} - \left[ \frac{2\pi\alpha a}{E^*} \right]^{1/2}. \quad (7)$$

Figure 5 is reproduced, with permission, from [98] (their Figure 5) in which they directly compare the behaviour of cells governed by (A) an explicit extended Hertzian model of adhesion (Section 3.2.1) with (B) the JKR theory model (Section 3.2.2). This study of the destabilisation of a monolayer shows clearly how the hysteresis effect between attachment and detachment of cells within the JKR model leads to fewer cells detaching from the substrate over the same timescale when compared with the extended Hertz model. For further details of the model parameters in these simulations, see [98].

For more details and simulation results of tumour growth under either the modified Hertzian or JKR adhesion forces see, for example, the references in Table 2.

#### 4. Additional aspects of the TM

This review will now consider selected modelling efforts of the mathematical and computational oncology community with regards to modelling tumour-TM interactions. In Section 4.1 cell-ECM interactions are discussed while in Section 4.3 cell-vessel interactions are considered.

##### 4.1. Tumour interactions with the ECM

The ECM, on a basic level, is composed of a structured mesh (matrix) of fibres (e.g. collagen and fibronectin) within a gel of glycoproteins. We have




Figure 5: Destabilization of a monolayer using the extended Hertz interaction (A) and the JKR-interaction energy (B). The numbers (I), (II), (III) denote the knocked-out control mechanisms which lead the destabilisation. (I) contact inhibition, (II) anchorage-dependent proliferation (III) anchorage-dependent apoptosis (anoikis). PERMISSION PENDING

297 previously discussed cell-cell adhesion but another important adhesive process  
 298 in cell biology is cell-matrix adhesion. Focal adhesions are protein complexes  
 299 which connect the cell's cytoskeleton to the ECM [9, 100]. Focal adhesions  
 300 not only directly and mechanically link the cell to the ECM but they also act  
 301 as points of signalling (mechanotransduction); transmitting information about  
 302 the mechanics of the extracellular environment to cells through biochemical sig-  
 303 nalling molecules. Focal adhesion mechanotransduction plays an important role  
 304 in regulating both the shape and migration of cells [9]. Specific focal adhesion  
 305 proteins which act as mechanotransducers are the ECM protein, fibronectin,  
 306 and cell-membrane receptor integrins. Fibronectin also binds to collagen fibres  
 307 in the ECM. Collagen fibres give structure to tissue but also, naturally, by  
 308 extension, to the TM.

309 The fibrous connective tissue of the ECM performs a wide variety of functions  
 310 within the healthy body. In terms of cancer, and within the TM, the structure of  
 311 the ECM and the interaction of cancer cells with individual fibres of the matrix  
 312 drives both cell proliferation and migration. ECM binding is implicated, for  
 313 example, in proliferative signalling; experimental data, backed up by *in silico*  
 314 models, have shown that border cells (those connected to the ECM) of a MCTS  
 315 are less proliferative than cells in the interior [46]. Moreover, malignant cells  
 316 activate the “integrin migration pathway” and crawl towards and along the  
 317 protein network of the ECM; migration through the protein network results  
 318 in the rearrangement of the ECM structure as cancer cells use the integrin  
 319 pathway to cut-off the fibres and re-orient the ECM [101, 102]. Cell migration  
 320 can happen as a collective process that presents in different ways depending  
 321 on the tumour type and the nearby environment leading to different migration  
 322 structures [15, 17]. The physical properties of the environment itself affects



323 tumour development and progression. It is widely known that cells prefer stiff  
 324 matrices to softer ones (*durotaxis*, [8]). Tumours themselves are known to be  
 325 stiffer than normal tissue [103]. Furthermore, it has been shown that stiff ECM  
 326 promotes tumour progression [104, 105]. On the other hand it has been shown  
 327 that tumour-repopulating cells (TRCs) are more proliferative in soft rather than  
 328 stiff environments [10]. To fully understand cancer development and local tissue  
 329 invasion it is important to model the ECM alongside the cancer cells. To model  
 330 the ECM it is natural to incorporate fibres as additional agents within an agent-  
 331 based model.

#### 332 4.2. Cell-fibre interactions

333 In [88] the ECM fibres are modelled using a force-based, individual-based  
 334 model. Single-cell experiments are carried out to determine the affect that the  
 335 cell’s environment (in this case a 2D substrate) has on its migration. By placing  
 336 a single cell in a domain segregated by substrates with different matrix stiff-  
 337 nesses [88] were able to reproduce the experimental results of [8] showing that  
 338 cells are drawn preferentially to stiffer matrices, hypothesising that it was the  
 339 lack of matrix reorientation by the cell that drives *durotaxis*. In a second ex-  
 340 periment they showed the observable “follow-the-leader” behaviour of collective  
 341 cell migration [106]. Figure 6 reproduces, with permission, their Figure 10, in  
 342 which a single non-polarised cell becomes polarised and “follows” the path of  
 343 polarised “leader” cell.

344 In [85] the 2D model of [88] is extended to 3D and matrix fibres are in-  
 345 corporated into a CBM for tumour growth. Each individual fibre is modelled  
 346 explicitly by a thin cylinder (described by its extrema and radius), and the  
 347 three-dimensional computational domain is filled with fibres of a given distri-  
 348 bution of positions and orientations. In a similar way to cell-cell interactions,  
 349 cell-fibre interactions are governed by attractive and repulsive forces; a cell in  
 350 contact with a fibre will feel an adhesive force, parallel to fibre orientation and  
 351 a repulsive force orthogonal to the fibre [107]. The cell-fibre interaction force is  
 352 computed as the sum of these orthogonal/repulsive and parallel/adhesive terms,  
 353  $\mathbf{F}_{i,f} = F_{\parallel} - F_{\perp}$ . The combined force  $\mathbf{F}_{i,f}$  is added to the right-hand side of the  
 354 equation of motion of each cell (Equation 1). We outline the chosen forms of  
 355 the forces given in [85] in the following Section.

##### 356 4.2.1. Cell-fibre forces

357 The cell-fibre adhesive force between a cell,  $i$ , and fibre,  $f$ , is modelled in  
 358 [85] by

$$\mathbf{F}_{\parallel} = \alpha_{\text{fibre}} \left( 1 - \frac{\|\mathbf{v}_i\|}{v_{\max}} \right) \left( \frac{|\mathbf{v}_i \cdot \mathbf{l}_f|}{\|\mathbf{v}_i\|} \right)^s \mathbf{l}_f. \quad (8)$$

359 It is directed along the normalised direction vector the fibre,  $\mathbf{l}_f$ , and depends on  
 360 the normalised scalar product between fibre direction and cell velocity (polarity),  
 361  $\mathbf{v}_i \cdot (\hat{\mathbf{x}}_i)$ . Thus, this force is maximised when a cell is already travelling parallel  
 362 to the fibre in question. Moreover, the force depends on an adhesion coefficient,  
 363  $\alpha_{\text{fibre}}$ , and on a threshold velocity,  $v_{\max}$ , which limits the pulling effect of fibres.




Figure 6: Snapshots in time indicating how two cells collectively migrate through the matrix. A non-polarized cell (red in plot **a**) becomes polarized (turning green) and then follows the path of the existing polarized cell (green in plot **a**). Reprinted from [88], Copyright (2012), with permission from Elsevier.

364 The additional parameter  $s > 0$  is used to model additional effects which might  
 365 increase ( $s < 1$ ) or decrease ( $s > 1$ ) the pulling effect.

366 The cell-fibre repulsion force is modelled via an additional friction exerted  
 367 by the fibre, given in [85] by

$$\mathbf{F}_{\perp} = \beta_{\text{fibre}} \left( \frac{\|\mathbf{v}_i\|^2 - |\mathbf{v}_i \cdot \mathbf{l}_f|^2}{\|\mathbf{v}_i\|^2} \right)^r \mathbf{v}_i. \quad (9)$$

368 It is directed parallel to cell velocity and depends on the component of cell  
 369 velocity orthogonal to the fibre, being maximised when the cell is travelling  
 370 directly orthogonal to the fibre in question. The coefficient of cell-fibre friction  
 371 is  $\beta_{\text{fibre}}$  and the exponent  $r > 0$  can be used to model nonlinear effects which  
 372 increase ( $r < 1$ ) or decrease ( $r > 1$ ) the repulsive forces.

373 Figure 7 is reproduced, with permission, from [85] (their Figure 4) shows how  
 374 a tumour develops oriented along fibres which are uniformly distributed aligned  
 375 with the  $y$ -axis. Initially a single cancer cell is placed within a fibrous domain,  
 376 the resulting tumour which has developed (after 9 000 timesteps, approximately  
 377 6 days) is shown in Figure 7. Whereas, in the absence of fibres, one would  
 378 typically see a spherical tumour mass form (as in Figure 4), here the growth  
 379 has been stretched out along the fibrous tissue. For further details of the model  
 380 and associated parameters, see [85].




Figure 7: Figures showing the results of a simulation of tumour growth within a domain of uniformly distributed fibres (aligned with the  $y$ -axis) after 9 000 time steps. Cells are represented by red spheres, fibres in grey. Left: View orthogonal to the fibre orientation ( $xz$ -plane). Right: View in the  $yz$ -plane, cropped on the left side. Reprinted from [85], Copyright (2020), with permission from Elsevier.

Figure 8 (simulated under the model of [85]) shows the migration of a single (non-proliferating) cell within a given fibrous domain. On the lefthand of the domain fibres are directed at 45 degrees to the  $x$ -axis while they are aligned parallel to the  $x$ -axis on the righthand of the domain. The cell is placed at (250,50,250) shown by the blue circle. The simulations is run 50 times for 10 000 timesteps (approximately 7 days), while the path of the cell through the fibrous domain is monitored. The trajectories of the cell for each simulation are indicated by the light grey lines, with the final position marked in red. The mean path is indicated with the dark grey line. As can be seen the cell paths follow the orientation of the fibres, switching alignment as they cross from the left to righthand of the domain.

A further biologically relevant aspect that links cancer cells to the ECM is matrix re-modelling. Matrix metalloproteinases (MMPs) are enzymes which degrade ECM proteins (e.g. collagen fibres) through proteolysis. Proteolytic re-modelling of the ECM by MMPs is a key step towards cancer invasion [6]. Fibre degradation is taken into account in current state-of-the-art continuum models, see, for example, [108]. Alternative models of cell-ECM interactions include [109] who use Hookean springs which act via the basement membrane which links cells to the connective tissue.

#### 4.3. Tumour interactions with the Vasculature

Another important aspect of the TM is the vasculature. Blood vessels weave through the tissue supplying it with oxygen and other vital nutrients. Cell-vessel interactions are both mechanical and biochemical.

##### 4.3.1. Mechanical cell-vessel interactions

Cells interact mechanically with segments of the vessel network. In [85] they assume that repulsive and adhesive forces act between a cell and a vessel




Figure 8: Cell migration simulation within a segregated domain of distributed fibres (not shown) after 10 000 time steps. The distribution of the fibres is indicated by the cartoons, being different on the lefthand and righthand sides. The initial position of the cell is indicated by the blue circle and the final positions (of 50 simulations) by red circles with paths shown in light grey. The average path is depicted in dark grey. Unpublished result from [85].

segment and that these forces are analogous to those between cells (Section 3),  
for further details see [85]. Their simulations show tumours developing and  
embedding within pre-existing vasculature. The proliferation of cancer cells  
around blood vessels - modelling so called “tumour cords” is simulated in [89].  
In the case of a tumour chord rather than a spherical tumour growing with the  
classical radial profile (necrotic core, quiescent and proliferative outer ring) the  
opposite profile is derived with necrotic regions on the outside furthest away  
from the central blood vessel(s). Figure 9 is reproduced, with permission, from  
[89] (their Figure 15).

In order for cancer to metastasise and spread to secondary sites around the  
body, cancer cells must be able to access the vessel network. Intravasation (and  
its analogous reverse, extravasation) is the process by which a cell enters (or  
leaves) the vascular network. In [92] they model the key metastatic process of  
intravasation using a CBM coupled to a deterministic model of the intracellular  
protein pathways which allow cells to migrate through the vessel endothelial  
wall (transendothelial migration, TEM) [110, 111]. In this case adhesion of the  
cancer cell with the vessel endothelia is key, and as before adhesion is driven by  
cadherins. Vascular endothelial cadherins (VE-cadherin) bind the cells of the  
vessel wall together. A cancer cell disrupts endothelial bonds binding itself to  
the wall using N-cadherin. Figure 10 is reproduced, with permission, from [92]  
and shows a single cell approach and then intravasate a vessel wall.




Figure 9: Simulation results of a tumour cord interacting with two blood vessels (black cells are necrotic). (a-b) Tumour cord growing around two vessels, (c) oxygen profile levels in the tumour cord, (d) cross-section showing corresponding development of tumour cells. Reprinted by permission from Springer Nature Customer Service Centre GmbH: Springer Nature, *Bulletin of Mathematical Biology*, [89], COPYRIGHT (2018).

#### 4.3.2. Biochemical interactions - The hypoxic phenotype

Cancer cells, like normal cells, respond to the availability of oxygen, although the malignant response is anything but normal. We can characterise cancer cells into phenotypes based on their access to oxygen (e.g. normoxic, hypoxic and necrotic). Hypoxic cells are chronically lacking in sufficient oxygen, this deficiency of the main cell nutrient rather than being tumour suppressing actually drives tumour progression in numerous ways [5]. Jain lists the following responses of tumour cells to hypoxia: switch to anaerobic metabolism; resist apoptosis; undergo the epithelial-mesenchymal transition (EMT); induce a cancer stem-cell “repopulating” phenotype, resist anti-cancer therapies; cause inflammation and immunosuppression; genomic instability and angiogenic. Notice that these classical behaviours are closely aligned with the *Hallmarks of Cancer* [1, 2]; the hypoxic phenotype is what drives cancer progression and makes it so deadly.

Hypoxia is a main driver of the epithelial-mesenchymal transition (EMT) [112]. The EMT occurs when epithelial cells detach (losing their cell-cell adhesion and polarity) and gain mesenchymal cell attributes (migration, invasion and differentiation). The EMT is the first step towards cancer metastasis. In [113] they model the EMT and metastasis using a hybrid on-lattice individual




Figure 10: Spatio-temporal evolution dynamics of a malignant cell (red nucleus coloured cell, marked with a full arrow) approaching a blood vessel to undergo TEM. When the malignant cell attaches to the vessel, the VE-cadherin bonds are disrupted and new N-cadherin bonds are formed (shown in yellow). After some time, the malignant cell manages to disrupt the endothelial bonds enough to open a gap in the vessel and undergo TEM. Reproduced with permission from [92]. PERMISSION PENDING.

447 based approach. Hypoxia also drives angiogenesis, with hypoxic tumour cells  
 448 releasing vascular-endothelial growth factor (VEGF) which signals for tumour  
 449 angiogenesis. McDougall and coworkers are leading experts in modelling an-  
 450 giogenesis [114–118]. In [84] they incorporate normoxic, hypoxic and necrotic  
 451 phenotypes into a CBM to show how the hypoxia phenotype is implicated in the  
 452 formation of pseudopalisades (hypercellular “walls” surrounding necrotic zones)  
 453 in glioblastoma.

## 454 5. Conclusions

455 This paper provides a selective review of *in silico* models for tumour growth  
 456 and development, with specific emphasis on centre-based force-based agent  
 457 based models. Key authors in the field include Drasdo and coworkers [86, 97,  
 458 98, 119–121] while a great many other authors are contributing to this vibrant  
 459 area of research [84, 85, 122–126]. For a critical evaluation of the available  
 460 agent based modelling techniques, their advantages and disadvantages see, for  
 461 example, [69]. No review of such models would be complete without mentioning  
 462 the work of Macklin and co-authors [127–129] who have recently launched *Physi-*  
 463 *Cell* a comprehensive open source C++ code designed to simulate the growth  
 464 of tumours within the TM [130]. One aspect of the TM which has not been  
 465 discussed here, although which is a vital part, are tumour-associated immune  
 466 cells. *PhysiCell* has been used to model how immune cells attack a MCTS [130],  
 467 other agent-based models of tumour immune interactions-include [131–134].

468 The main take home message is that biomechanics need to be taken into  
 469 account. One might contrast individual-based models with reaction-diffusion

models of cancer. While reaction-diffusion models (for example, [135–137]) do offer insight they do not include biomechanics nor can they account for phenotypic variations that are well captured through an agent-based force-based approach. Even for the subset of reaction-diffusion-taxis models ([138], for example) where biomechanics may be implied they are not taken into account explicitly. Individual-based modelling, then, has significant advantages over reaction-diffusion models in determining the key mechanisms which drive metastatic spread. Perhaps in the future effort should be put into integrating reaction-diffusion models with biomechanics in order to gain the advantages of both approaches.

Agent-based modelling of tumour growth, however, is just a single strategy in the global effort of the scientific community in the fight against cancer. Indeed mathematical (and computational) oncology is a growing field in which research is being done on a broad range of topics spanning from modelling intracellular genetic pathways (see, for example, [139–141]) to modelling cancer therapies (see, for example, [142–144]). Looking to the future a multi-scale model of a growing tumour within the TM should seek to bring together not only the biomechanical aspects laid out above but equally other aspects from the diverse field of study. By incorporating intracellular pathways (such as in [91, 93]) which results in phenotypic differences between cells it is possible to derive a realistic heterogeneous cancer cell population. By using imaging combined with the modelling techniques above to render *in vivo* tumours *in silico* it is possible to simulate in real time and space the development of tumours predicting how they will invade and metastasise. By trialing cancer therapies on *in silico* tumours (as in [47, 118, 145]) clinicians can devise optimal therapy protocols that can at once become both the standard of care and patient specific. In combination these techniques will truly push the frontier of our understanding of cancer and lead towards personalised medicine where each patient can be treated truly individually.

## Acknowledgements

CKM gratefully acknowledges the support of EPSRC Grant No. EP/N014642/1 (EPSRC Centre for Multiscale Soft Tissue Mechanics - With Application to Heart & Cancer).

## References

- [1] D. Hanahan and R. A. Weinberg. Hallmarks of cancer. Cell, 100:57–70, 2000.
- [2] D. Hanahan and R. A. Weinberg. Hallmarks of cancer: the next generation. Cell, 144:646–674, 2011.
- [3] Y. Kim, M. A. Stolarska, and H. G. Othmer. The role of the microenvironment in tumor growth and invasion. Prog Biophys Mol Biol, 106:353–379, 2011.

511 [4] G. D. Yancopoulos, S. Davis, N. W. Gale, J. S. Rudge, S. J. Wiegand,  
512 and J. Holash. Vascular-specific growth factors and blood vessel formation.  
513 Nature, 407(6801):242–248, 2000.

514 [5] R. K. Jain. Antiangiogenesis strategies revisited: from starving tumors to  
515 alleviating hypoxia. Cancer Cell, 26(5):605–622, 2014.

516 [6] L. M. Coussens and Z. Werb. Matrix metalloproteinases and the develop-  
517 ment of cancer. Chem Biol, 3(11):895–904, 1996.

518 [7] Y. Itoh and H. Nagase. Matrix metalloproteinases in cancer. Essays  
519 Biochem, 38:21–36, 2000.

520 [8] C.-M. Lo, H.-B. Wang, M. Dembo, and Y.-L. Wang. Cell movement is  
521 guided by the rigidity of the substrate. Biophys J, 79(1):144–152, 2000.

522 [9] J. Seong, N. Wang, and Y. Wang. Mechanotransduction at focal adhe-  
523 sions: from physiology to cancer development. J Cell Mol Med, 17(5):597–  
524 604, 2013.

525 [10] J. Liu, Y. Tan, H. Zhang, Y. Zhang, P. Xu, J. Chen, Y. C. Poh, K. Tang,  
526 N. Wang, and B. Huang. Soft fibrin gels promote selection and growth of  
527 tumorigenic cells. Nat Mater, 11(8):734–741, 2012.

528 [11] T. Brabletz, A. Jung, S. Reu, M. Porzner, F. Hlubek, L. A. Kunz-  
529 Schughart, R. Knuechel, and T. Kirchner. Variable beta-catenin expres-  
530 sion in colorectal cancers indicates tumor progression driven by the tumor  
531 environment. Proc Natl Acad Sci USA, 98:10356–10361, 2001.

532 [12] T. Brabletz, S. Spaderna, J. Kolb, F. Hlubek, G. Faller, C. J. Bruns,  
533 A. Jung, J. Nentwich, I. Duluc, C. Domon-Dell, T. Kirchner, and J. N.  
534 Freund. Down-regulation of the homeodomain factor cdx2 in colorectal  
535 cancer by collagen type i: an active role for the tumor environment in  
536 malignant tumor progression. Cancer Res, 64:6973–6977, 2004.

537 [13] H. Dillekås, M. S. Rogers, and O. Straume. Are 90% of deaths from cancer  
538 caused by metastases? Cancer Med, 8:5574–5576, 2019.

539 [14] P. Friedl, K. S. Zänker, and E.-B. Bröcker. Cell migration strategies in 3-d  
540 extracellular matrix: Differences in morphology, cell matrix interactions,  
541 and integrin function. Microsc Res Tech, 43:369–378, 1998.

542 [15] P. Fiedl and K. Wolf. Tumour-cell invasion and migration: diversity and  
543 escape mechanisms. Nat Rev Cancer, 3(5):362–374, 2003.

544 [16] K. Wolf, Y. I. Wu, Y. Liu, J. Geiger, E. Tam, C. Overall, M. S. Stack, and  
545 P. Friedl. Multi-step pericellular proteolysis controls the transition from  
546 individual to collective cancer cell invasion. Nat Cell Biol, 9(8):893–904,  
547 2007.



- 548 [17] P. Fiedl and K. Wolf. Proteolytic interstitial cell migration: a five-step  
549 process. Cancer Metastasis Rev, 28(1-2):129–135, 2009.
- 550 [18] P. Fiedl and S. Alexander. Cancer invasion and the microenvironment:  
551 plasticity and reciprocity. Cell, 147(5):992–1009, 2011.
- 552 [19] J. A. Davies, editor. Chapter 11 - Guidance by Contact, pages 129–145.  
553 Academic Press, 2013.
- 554 [20] T. Iskratsch, H. Wolfenson, and M. P. Sheetz. Appreciating force and  
555 shape the rise of mechanotransduction in cell biology. Nat Rev Mol Cell  
556 Biol, 15:825–833, 2014.
- 557 [21] L. Chin, Y. Xia, D. E. Discher, and P. A. Janmey. Mechanotransduction  
558 in cancer. Curr Opin Chem Eng, 11:77–84, 2016.
- 559 [22] G. Helmlinger, P. A. Netti, H. C. Lichtenbeld, R. J. Melder, and R. K.  
560 Jain. Solid stress inhibits the growth of multicellular tumor spheroids.  
561 Nat Biotechnol, 15:778–783, 1997.
- 562 [23] G. Cheng, J. Tse, R. K. Jain, and L. L. Munn. Micro-environmental mech-  
563 anical stress controls tumor spheroid size and morphology by suppressing  
564 proliferation and inducing apoptosis in cancer cells. PLoS ONE, 4:e4632,  
565 2009.
- 566 [24] J. Yang, S. A. Mani, J. L. Donaher, S. Ramaswamy, C. Itzykson, R.  
567 A. Come, P. Savagner, I. Gitelman, A. Richardson, and Weinberg R. A.  
568 Twist, a master regulator of morphogenesis, plays an essential role in  
569 tumor metastasis. Cell, 117(7):927–939, 2004.
- 570 [25] M. Basan, T. Risler, J. F. Joanny, X. S. Garau, and J. Prost. Homeostatic  
571 competition drives tumor growth and metastasis nucleation. HFSP J,  
572 3(4):265–272, 2009.
- 573 [26] D. T. Butcher, T. Alliston, and V. M. Weaver. A tense situation: Forcing  
574 tumour progression. Nat Rev Cancer, 9:108–122, 2009.
- 575 [27] J. M. Tse, G. Cheng, J. A. Tyrrell, S. A. Wilcox-Adelman, Y. Boucher,  
576 R. K. Jain, and L. L. Munn. Mechanical compression drives cancer cells  
577 toward invasive phenotype. PNAS, 109(3):911–916, 2012.
- 578 [28] H. P. Greenspan. Models for the growth of a solid tumour by diffusion.  
579 Stud Appl Math, 52:317–340, 1972.
- 580 [29] U. Del Monte. Does the cell number 10(9) still really fit one gram of tumor  
581 tissue? Cell Cycle, 8(3):505–506, 2009.
- 582 [30] R. P. Araujo and McElwain D. L. S. A history of the study of solid  
583 tumour growth: the contribution of mathematical modelling. Bull Math  
584 Biol, 66(5):1039–1091, 2004.

- 585 [31] T. Roose, S. J. Chapman, and P. K. Maini. Mathematical models of  
586 avascular tumor growth. SIAM Rev, 49(2):179–208, 2007.
- 587 [32] A. F. Jones, H. M. Byrne, J. S. Gibson, and J. W. Dold. A mathematical  
588 model of the stress induced during avascular tumour growth. J Math Bio,  
589 40(6):473–499, 2000.
- 590 [33] D. Ambrosi and F. Mollica. On the mechanics of a growing tumor. Int J  
591 Eng Sci, 40(12):1297–1316, 2002.
- 592 [34] C. Koike, T. D. McKee, A. Pluen, S. Ramanujan, K. Burton, L. L. Munn,  
593 Y. Boucher, and R. K. Jain. Solid stress facilitates spheroid formation:  
594 potential involvement of hyaluronan. Br J Cancer, 86:947–953, 2002.
- 595 [35] H. Byrne and L. Preziosi. Modelling solid tumour growth using the theory  
596 of mixtures. Math Med Biol, 20:341–366, 2003.
- 597 [36] D. Ambrosi and F. Mollica. The role of stress in the growth of a multicell  
598 spheroid. J Math Biol, 48(5):477–479, 2004.
- 599 [37] H. Byrne and M. A. J. Chaplain. Modelling the role of cell-cell adhesion  
600 in the growth and development of carcinomas. Math Comput Model,  
601 12:1–17, 1996.
- 602 [38] H. Byrne. The importance of intercellular adhesion in the development of  
603 carcinomas. IMA J Math Appl Med Biol, 14:305–323, 1997.
- 604 [39] C. Breward, H. Byrne, and C. Lewis. The role of cell-cell interactions in a  
605 two-phase model for avascular tumour growth. J Math Biol, 45:125–152,  
606 2002.
- 607 [40] H. M. Byrne, J. R. King, D. L. S. McElwain, and L. Preziosi. A two-phase  
608 model of solid tumor growth. Appl Math Lett, 16(4):567–573, 2003.
- 609 [41] N. J. Armstrong, K. J. Painter, and J. A. Sherratt. A continuum approach  
610 to modelling cell-cell adhesion. J Theor Biol, 243:98–113, 2006.
- 611 [42] M. A. J. Chaplain, L. Graziano, and L. Preziosi. Mathematical modelling  
612 of the loss of tissue compression responsiveness and its role in solid tumour  
613 development. Math Med Biol, 23:197–229, 2006.
- 614 [43] I. Ramis-Conde, M. A. J. Chaplain, and A. Anderson. Mathematical  
615 modelling of cancer cell invasion of tissue. Math Comput Model, 47:533–  
616 545, 2008.
- 617 [44] L. Preziosi and A. Tosin. Multiphase modeling of tumor growth and  
618 extracellular matrix interaction: mathematical tools and applications. J  
619 Math Biol, 58(4-5):625–656, 2008.

- [45] A. R. Kansal, S. Torquato, G. R. IV Harsh, E. A. Chiocca, and T. S. Deisboeck. Simulated brain tumor growth dynamics using a three dimensional cellular automaton. J Theor Biol, 203(4):367–382, 2000.
- [46] N. Jagiella, B. Müller, M. Müller, I. E. Vignon-Clementel, and D. Drasdo. Inferring growth control mechanisms in growing multi-cellular spheroids of nscic cells from spatial-temporal image data. PLoS Comput Biol, 12(2):e1004412, 2016.
- [47] S. C. Brüningk, P. Ziegenhein, I. Rivens, U. Oelfke, and G. T. Haar. A cellular automaton model for spheroid response to radiation and hyperthermia treatments. Sci Rep, 9(1):17674, 2019.
- [48] A. A. Patel, E. T. Gawlinski, S. K. Lemieux, and R. A. Gatenby. A cellular automaton model of early tumor growth and invasion: The effects of native tissue vascularity and increased anaerobic tumor metabolism. J Theor Biol, 213(3):315–331, 2001.
- [49] A. R. A. Anderson. A hybrid model of solid tumour invasion: the importance of cell adhesion. Math Med Biol, 22:163186, 2006.
- [50] A. R. A. Anderson, A. M. Weaver, P. T. Cummings, and V. Quaranta. Tumor morphology and phenotypic evolution driven by selective pressure from the microenvironment. Cell, 17(5):905–915, 2006.
- [51] M. Aubert, M. Badoual, S. Freol, C. Christov, and B. Grammaticos. A cellular automaton model for the migration of glioma cells. Phys Biol, 3:93–100, 2006.
- [52] J. Vivas, D. Garzón-Alvarado, and M. Cerrolaza. Modeling cell adhesion and proliferation: a cellular-automata based approach. Adv Model Simul Eng Sci, 2:32, 2015.
- [53] M. Block, E. Schöll, and D. Drasdo. Classifying the expansion kinetics and critical surface dynamics of growing cell populations. Phys Rev Lett, 99(24):248101, 2007.
- [54] A. R. A. Anderson, K. A. Rejniak, P. Gerlee, and V. Quaranta. Modelling of cancer growth, evolution and invasion: Bridging scales and models. Math Model Nat Phenom, 2(3):1–29, 2007.
- [55] L. Zhang, C. G. Strouthos, Z. Wang, and T. S. Deisboeck. Simulating brain tumor heterogeneity with a multiscale agent-based model: Linking molecular signatures, phenotypes and expansion rate. Math Comput Model, 49(1-2):307–319, 2009.
- [56] H. Enderling and P. Hahnfeldt. Cancer stem cells in solid tumors: Is ‘evading apoptosis’ a hallmark of cancer? Prog Biophys Mol Biol, 106(2):391–399, 2011.

- [57] Y. Cai, J. Wu, S. Xu, and Z. Li. A hybrid cellular automata model of multicellular tumour spheroid growth in hypoxic microenvironment. J Appl Math, 2013:519895, 2013.
- [58] M. Radszuweit, M. Block, J. Hengstler, E. Schöll, and D. Drasdo. Comparing the growth kinetics of cell populations in two and three dimensions. Phys Rev E, 79:051907, 2009.
- [59] D. Chen, Y. Jiao, and S. Torquato. A cellular automaton model for tumor dormancy: emergence of a proliferative switch. PloS one, 9(10):e109934, 2014.
- [60] E. Stott, N. Britton, J. Glazier, and M. Zajac. Stochastic simulation of benign avascular tumor growth using the potts model. Math Comput Model, 30:183–198, 1999.
- [61] S. Turner and J.A. Sherratt. Intercellular adhesion and cancer invasion: a discrete simulation using the extended potts model. J Theor Biol, 216:85–100, 2002.
- [62] E. Boghaert, D. C. Radisky, and C. M. Nelson. Lattice-based model of ductal carcinoma in situ suggests rules for breast cancer progression to an invasive state. PLoS Comput Biol, 10:e1003997, 2014.
- [63] A. Shirinifard, J. S. Gens, B. L. Zaitlen, N. J. Popawski, M. Swat, and J. A. Glazier. 3d multi-cell simulation of tumor growth and angiogenesis. PLoS ONE, 4(10):e7190, 2009.
- [64] B. M. Rubenstein and L. J. Kaufman. The role of extracellular matrix in glioma invasion: a cellular potts model approach. Biophys J, 95:5661–5680, 2008.
- [65] J. F. Li and J. Lowengrub. The effects of cell compressibility, motility and contact inhibition on the growth of tumor cell clusters using the cellular potts model. J Theor Biol, 343:79–91, 2014.
- [66] S. Dormann and A. Deutsch. Modeling of self-organized avascular tumor growth with a hybrid cellular automaton. In Silico Biol, 2:393–406, 2002.
- [67] M. Wurzel, C. Schaller, M. Simon, and A. Deutsch. Cancer cell invasion of normal brain tissue: Guided by prepattern? J Theor Med, 6(1):21–31, 2005.
- [68] H. Hatzikirou and A. Deutsch. Cellular automata as microscopic models of cell migration in heterogeneous environments. Curr Top Dev Biol, 81:401–434, 2008.
- [69] P. Van Liedekerke, M. M. Palm, N. Jagiella, and D. Drasdo. Simulating tissue mechanics with agent-based models: concepts, perspectives and some novel results. Comp Part Mech, 2:401–444, 2015.

- [70] J. Moreira and A. Deutsch. Cellular automaton models of tumor development: a critical review. Adv Complex Syst, 5:247–268, 2002.
- [71] M. S. Alber, M. A. Kiskowski, J. A. Glazier, and Y. Jiang. On cellular automaton approaches to modeling biological cells. In J. Rosenthal and D. S. Gilliam, editors, Mathematical systems theory in biology, communication, and finance, pages 1–39. The IMA Volumes in Mathematics and its Applications, vol 134. Springer, New York, NY, 2003.
- [72] A. R. A. Anderson, M. A. J. Chaplain, and K. A. Rejniak. Single-cell-based models in biology and medicine. Birkhäuser, Basel, 2007.
- [73] A. Szabó and R. M. H. Merks. Cellular pots modeling of tumor growth, tumor invasion and tumor evolution. Front Oncol, 3:87, 2013.
- [74] H. Hatzikirou, G. Breier, and A. Deutsch. Cellular automaton modeling of tumor invasion. In R. Meyers, editor, Encyclopedia of Complexity and Systems Science, pages 1–13. Springer, Berlin, Heidelberg, 2014.
- [75] J. Metzcar, Y. Wang, R. Heiland, and P. Macklin. A review of cell-based computational modeling in cancer biology. JCO Clin Cancer Inform, 3:1–13, 2019.
- [76] N. Ouchi, J. A. Glazier, J. Rieu, A. Upadhyaya, and Y. Sawada. Improving the realism of the cellular pots model in simulations of biological cells. Physica A, 329(3-4):451–458, 2003.
- [77] M. Scianna and L. Preziosi. A hybrid model describing different morphologies of tumor invasion fronts. Math Model Nat Phenom, 7(1):78–104, 2012.
- [78] J. T. Daub and R. M. H. Merks. A cell-based model of extracellularmatrix-guided endothelial cell migration during angiogenesis. Bull Math Biol, 75(8):1377–1399, 2013.
- [79] M. H. Swat, G. L. Thomas, J. M. Belmonte, A. Shirinifard, D. Hmeljak, and J. A. Glazier. Chapter 13 - multi-scale modeling of tissues using compucell3d. In A. R. Asthagiri and A. P. Arkin, editors, Computational Methods in Cell Biology, volume 110 of Methods in Cell Biology, pages 325–366. Academic Press, 2012.
- [80] K. A. Rejniak. A single-cell approach in modeling the dynamics of tumor microregions. Math Biosci Eng, 2:643–655, 2005.
- [81] K. A. Rejniak. An immersed boundary framework for modelling the growth of individual cells: an application to the early tumour development. J Theor Biol, 247:186–204, 2007.
- [82] K. A. Rejniak and R. H. Dillon. A single cell-based model of the ductal tumour microarchitecture. Comput Math Methods Med, 8(1):51–69, 2007.

- [83] K. A. Rejniak, S. E. Wang, N. S. Bryce, H. Chang, B. Parvin, J. Jourquin, L. Estrada, J. W. Gray, C. L. Arteaga, A. M. Weaver, V. Quaranta, and A. R. A. Anderson. Linking changes in epithelial morphogenesis to cancer mutations using computational modeling. *PLoS Comput Biol*, 6(8):e1000900, 2010.
- [84] A. Caiazzo and I. Ramis-Conde. Multiscale modelling of palisade formation in glioblastoma multiforme. *J Theor Biol*, 383:145–156, 2015.
- [85] C. K. Macnamara, A. Caiazzo, I. Ramis-Conde, and M. A. J. Chaplain. Computational modelling and simulation of cancer growth and migration within a 3d heterogeneous tissue: the effects of fibre and vascular structure. *J Comp Sci*, 40:101067, 2020.
- [86] D. Drasdo and S. Hoehme. Individual-based approaches to birth and death in avascular tumors. *Math Comput Model*, 37:1163–1175, 2003.
- [87] H. Byrne and D. Drasdo. Individual-based and continuum models of growing cell populations: a comparison. *J Math Biol*, 58(657):301–313, 2009.
- [88] D. K. Schlüter, I. Ramis-Conde, and M. A. J. Chaplain. Computational modeling of single-cell migration: the leading role of extracellular matrix fibers. *Biophys J*, 103:1141–1151, 2012.
- [89] Z. Szymańska, M. Cytowski, E. Mitchell, C. K. Macnamara, and M. A. J. Chaplain. Computational modelling of cancer development and growth: Modelling at multiple scales and multiscale modelling. *Bull Math Biol*, 80:1366–1403, 2018.
- [90] R. E. Mahaffy, S. Park, E. Gerde, J. Käs, and C. K. Shih. Quantitative analysis of the viscoelastic properties of thin regions of fibroblasts using atomic force microscopy. *Biophys J*, 86(3):1777–1793, 2004.
- [91] I. Ramis-Conde, D. Drasdo, A. R. A. Anderson, and M. A. J. Chaplain. Modeling the influence of the e-cadherin-beta-catenin pathway in cancer cell invasion: a multiscale approach. *Biophys J*, 95(1):155–65, 2008.
- [92] I. Ramis-Conde, M. A. J. Chaplain, A. R. A. Anderson, and D. Drasdo. Multi-scale modelling of cancer cell intravasation: the role of cadherins in metastasis. *Phys Biol*, 6(1):016008, 2009.
- [93] I. Ramis-Conde and D. Drasdo. From genotypes to phenotypes: classification of the tumour profiles for different variants of the cadherin adhesion pathway. *Phys Biol*, 9(3):036008, 2012.
- [94] D. K. Schlüter, I. Ramis-Conde, and M. A. J. Chaplain. Multi-scale modelling of the dynamics of cell colonies: insights into cell-adhesion forces and cancer invasion from in silico simulations. *J R Soc Interface*, 12:20141080, 2015.

- 772 [95] H. Hertz. Ueber die berührung fester elastischer körper (on the contact  
773 of elastic solids). J Reine Angew Mat, 92:156–171, 1882.
- 774 [96] J. Galle, M. Loeffler, and D. Drasdo. Modeling the effect of deregulated  
775 proliferation and apoptosis on the growth dynamics of epithelial cell pop-  
776 ulations in vitro. Biophys J, 88(1):62–75, 2005.
- 777 [97] D. Drasdo and S. Hoehme. A single-cell-based model of tumor growth in  
778 vitro: monolayers and spheroids. Phys biol, 2:133–47, 2005.
- 779 [98] D. Drasdo, S. Hoehme, and M. Block. On the role of physics in the growth  
780 and pattern formation of multicellular systems: What can we learn from  
781 individual-cell based models? J Statist Phys, 128:287–345, 2007.
- 782 [99] K. L. Johnson, K. Kendall, and A. Roberts. Surface energy and the contact  
783 of elastic solids. Proceedings of the Royal Society of London. Series A,  
784 Mathematical and Physical Sciences, 324(1558):301–313, 1971.
- 785 [100] W. H. Goldmann. Mechanotransduction and focal adhesions. Cell Biol  
786 Int, 36(7):649–652, 2012.
- 787 [101] J. Hood and D. Cheresch. Role of integrins in cell invasion and migration.  
788 Nat Rev Cancer, 2:91–100, 2002.
- 789 [102] M. W. Pickup, J. K. Mouw, and V. M. Weaver. The extracellular matrix  
790 modulates the hallmarks of cancer. EMBO Rep, 15:1243–1253, 2014.
- 791 [103] A. Nagelkerke, J. Bussink, A. E. Rowan, and P. N. Span. The mechanical  
792 microenvironment in cancer: How physics affects tumours. Semin Cancer  
793 Biol, 35:62–70, 2015.
- 794 [104] K. R. Levental, H. Yu, L. Kass, J. N. Lakins, M. Egeblad, J. T. Erler,  
795 S. F. Fong, K. Csiszar, A. Giaccia, W. Weninger, M. Yamauchi, D. L.  
796 Gasser, and V. M. Weaver. Matrix crosslinking forces tumor progression  
797 by enhancing integrin signaling. Cell, 139(5):891–906, 2009.
- 798 [105] P. P. Provenzano, D. R. Inman, K. W. Eliceiri, and P. J. Keely. Matrix  
799 density-induced mechanoregulation of breast cell phenotype, signaling and  
800 gene expression through a fak-erk linkage. Oncogene, 28(49):4326–4343,  
801 2009.
- 802 [106] P. Friedl and K. Wolf. Plasticity of cell migration: a multiscale tuning  
803 model. J Cell Biol, 188:11–19, 2010.
- 804 [107] J. C. Dallan, J. A. Sherratt, and P. K. Maini. Mathematical modelling of  
805 extracellular matrix dynamics using discrete cells: fiber orientation and  
806 tissue regeneration. J Theor Biol, 199:449–471, 1999.
- 807 [108] R. Shuttleworth and D. Trucu. Multiscale modelling of fibres dynamics  
808 and cell adhesion within moving boundary cancer invasion. Bull Math  
809 Biol, 81:2176–2219, 2019.

- [109] G. D’Antonio, P. Macklin, and L. Preziosi. An agent-based model for elasto-plastic mechanical interactions between cells, basement membrane and extracellular matrix. Math Biosci Eng, 10(1):75–101, 2013.
- [110] S. P. Chiang, R. M. Cabrera, and J. E. Segall. Tumor cell intravasation. Am J Physiol Cell Physiol, 311(1):C1–C14, 2016.
- [111] M. V. Zavyalova, E. V. Denisov, L. A. Tashireva, O. E. Savelieva, E. V. Kaigorodova, N. V. Krakhmal, and V. M. Perelmuter. Intravasation as a key step in cancer metastasis. Biochemistry (Mosc), 84(7):762–772, 2019.
- [112] L. Zhang, G. Huang, X. Li, Y. Zhang, Y. Jiang, J. Shen, J. Liu, Q. Wang, J. Zhu, X. Feng, J. Dong, and C. Qian. Hypoxia induces epithelial-mesenchymal transition via activation of *snail* by hypoxia-inducible factor-1 $\alpha$  in hepatocellular carcinoma. BMC Cancer, 13:108, 2013.
- [113] L. C. Franssen, T. Lorenzi, A. F. Burgess, and M. A. J. Chaplain. A mathematical framework for modelling the metastatic spread of cancer. Bull Math Biol, 81(6):1965–2010, 2019.
- [114] S. R. McDougall, A. R. Anderson, M. A. J. Chaplain, and J. A. Sherratt. Mathematical modelling of flow through vascular networks: implications for tumour-induced angiogenesis and chemotherapy strategies. Bull Math Biol, 64:673–702, 2002.
- [115] S. R. McDougall, A. R. Anderson, and M. A. J. Chaplain. Mathematical modelling of dynamic adaptive tumour-induced angiogenesis: clinical implications and therapeutic targeting strategies. J Theor Biol, 241:564–589, 2006.
- [116] M. J. Machado, M. G. Watson, A. H. Devlin, M. A. J. Chaplain, S. R. McDougall, and C. A. Mitchell. Dynamics of angiogenesis during wound healing: a coupled in vivo and in silico study. Microcirculation, 8:183–197, 2011.
- [117] M. G. Watson, S. R. McDougall, M. A. J. Chaplain, A. H. Devlin, and C. A. Mitchell. Dynamics of angiogenesis during murine retinal development: a coupled in vivo and in silico study. J. R. Soc. Interface, 9:2351–2364, 2011.
- [118] A. Boujelben, M. G. Watson, S. R. McDougall, Y.-F. Yen, E. R. Gerstner, C. Catana, T. Deisboeck, T. T. Batchelor, D. Boas, B. Rosen, J. Kalpathy-Cramer, and M. A. J. Chaplain. Multimodality imaging and mathematical modelling of drug delivery to glioblastomas. Interface Focus, 6:20160039, 2016.
- [119] D. Drasdo, R. Kree, and J. McCaskill. Monte-carlo approach to tissue-cell populations. Phys Rev E, 52(6):6635–6657, 1995.



- [120] J. Galle, G. Aust, G. Schaller, T. Beyer, and D. Drasdo. Individual cell-based models of the spatial-temporal organization of multicellular systems - achievements and limitations. Cytom Part A, 69:704–10, 2006.
- [121] P. Van Liedekerke, A. Buttenschön, and D. Drasdo. Off-lattice agent-based models for cell and tumor growth: numerical methods, implementation, and applications. In M. Cerrolaza, S. Shefelbine, and D. Garzón-Alvarado, editors, Numerical methods and advanced simulation in biomechanics and biological processes, pages 245–267. London, UK; San Diego, CA; Cambridge, MA; Oxford, UK: Elsevier Academic Press, 2018.
- [122] G. Schaller and M. Meyer-Hermann. Multicellular tumor spheroid in an off-lattice voronoi-delaunay cell model. Phys Rev E, 71:051910, 2019.
- [123] J. Galle, D. Sittig, I. Hanisch, M. Wobus, E. Wandel, M. Loeffler, and G. Aust. Individual cell-based models of tumor-environment interactions: Multiple effects of cd97 on tumor invasion. Am J Pathol, 169(5):1802–1811, 2006.
- [124] M. Cytowski and Z. Szymańska. Large scale parallel simulations of 3-d cell colony dynamics. IEEE Comput Sci Eng, 16(5):86–95, 2014.
- [125] M. Cytowski and Z. Szymańska. Enabling large scale individual-based modelling through high performance computing. In ITM Web of Conferences, volume 5, page 00014, 2015.
- [126] M. Cytowski and Z. Szymańska. Large scale parallel simulations of 3-d cell colony dynamics. ii. coupling with continuous description of cellular environment. Comput Sci Eng, 17:44–48, 2015.
- [127] P. Macklin and M. E. Edgerton. Agent-based cell modeling: application to breast cancer. In V. Cristini and J. S. Lowengrub, editors, Multiscale modeling of cancer: an integrated experimental and mathematical modeling approach, pages 206–234. Cambridge University Press, Cambridge, UK, 2010.
- [128] P. Macklin, M. E. Edgerton, A. Thompson, and V. Cristini. Patient-calibrated agent-based modelling of ductal carcinoma in situ (dcis): from microscopic measurements to macroscopic predictions of clinical progression. J Theor Biol, 301:122–170, 2012.
- [129] P. Macklin, S. Mumenthaler, and J. Lowengrub. Modeling multiscale necrotic and calcified tissue biomechanics in cancer patients: application to ductal carcinoma in situ (dcis). In Multiscale Computer Modeling in Biomechanics and Biomedical Engineering. Studies in Mechanobiology, Tissue Engineering and Biomaterials, volume 14, pages 349–380. Springer, Berlin, Heidelberg, 2013.

- [130] A. Ghaffarizadeh, R. Heiland, S. H. Friedman, S. M. Mumenthaler, and P. Macklin. Physicell: An open source physics-based cell simulator for 3-d multicellular systems. PLoS Comput Biol, 14(2):e1005991, 2018.
- [131] F. Pappalardo, S. Musumeci, and S. Motta. Modeling immune system control of atherogenesis. Bioinformatics, 24:1715–1721, 2008.
- [132] W.-Y. Hu, W.-R. Zhong, F.-H. Wang, L. Li, and Y.-Z. Shao. In silico synergism and antagonism of an anti-tumour system intervened by coupling immunotherapy and chemotherapy: a mathematical modelling approach. Bull Math Biol, 74(2):434–452, 2012.
- [133] J. N. Kather, J. Poleszczuk, M. Suarez-Carmona, J. Krisam, P. Charoentong, N. A. Valous, C. A. Weis, L. Tavernar, F. Leiss, E. Herpel, F. Klupp, A. Ulrich, M. Schneider, A. Marx, D. Jäger, and N. Halama. In silico modeling of immunotherapy and stroma-targeting therapies in human colorectal cancer. Cancer Res, 77(22):6442–6452, 2017.
- [134] F. R. Macfarlane, T. Lorenzi, and M. A. J. Chaplain. Modelling the immune response to cancer: an individual-based approach accounting for the difference in movement between inactive and activated t cells. Bull Math Biol, 80:1539–1562, 2018.
- [135] R. A. Gatenby and E. T. Gawlinski. A reaction-diffusion model of cancer invasion. Cancer Res., 56:5745–5753, 1996.
- [136] A. J. Perumpanani, J. A. Sherratt, J. Norbury, and H. M. Byrne. Biological inferences from a mathematical model for malignant invasion. Invasion Metastasis, 16:209–221, 1996.
- [137] J. A. Sherratt and M. A. J. Chaplain. A new mathematical model for avascular tumour growth. J Math Biol, 43:291–312, 2001.
- [138] A. Gerisch and M. A. J. Chaplain. Mathematical modelling of cancer cell invasion of tissue: Local and non-local models and the effect of adhesion. J Theor Biol, 505(4):684–704, 2008.
- [139] N. A. M. Monk. Oscillatory expression of *hes1*, *p53*, and *nf- $\kappa$ b* driven by transcriptional time delays. Curr Biol, 13:1409–1413, 2003.
- [140] M. Sturrock, A. J. Terry, D. P. Xirodimas, A. M. Thompson, and M. A. J. Chaplain. Spatio-temporal modelling of the *hes1* and *p53*-*mdm2* intracellular signalling pathways. J Theor Biol, 273:15–31, 2011.
- [141] J. Eliaš, L. Dimitrio, J. Clairambault, and R. Natalini. Modelling *p53* dynamics in single cells: physiologically based ode and reaction-diffusion pde models. Phys Biol, 11:045001, 2014.
- [142] R. Rockne, E. C. Jr Alvord, J. K. Rockhill, and K. R. Swanson. A mathematical model for brain tumor response to radiation therapy. J Math Biol, 58(4-5):561–578, 2009.

- 925 [143] D. Wodarz. Computational modeling approaches to studying the dynam-  
926 ics of oncolytic viruses. Math Biosci Eng, 10(3):939–957, 2013.
- 927 [144] H. Enderling, J. C. L. Alfonso, E. Moros, J. J. Caudell, and L. B. Harrison.  
928 Integrating mathematical modeling into the roadmap for personalized ad-  
929 aptive radiation therapy. Trends Cancer, 5(8):467–474, 2019.
- 930 [145] A. L. Jenner, F. Frascoli, A. C. F. Coster, and P. S. Kim. Enhancing  
931 oncolytic virotherapy: Observations from a voronoi cell-based model. J  
932 Theor Biol, 485:110052, 2020.